

## Ptinidae of Greece (Coleoptera: Bostrichoidea) - a commented checklist with descriptions of new species

Petr ZAHRADNÍK

Forestry and Game Management Research Institute  
Strnady 136, CZ-156 00 Praha - 5, Zbraslav, Czech Republic  
e-mail: zahradnik@vulhm.cz

### Taxonomy, new species, faunistics, Coleoptera, Ptinidae, Palaearctic Region, Greece

**Abstract.** A review of the fauna of Ptinidae (Coleoptera: Bostrichoidea) of Greece is presented. One hundred and fifty-four species have been known from Greece. Five new species are described here - *Episernus graecus* sp. nov., *Ernobius rebli* sp. nov., *Xyletinus (Xyletinus) konvickai* sp. nov., *Xyletinus (Xyletinus) siteki* sp. nov. and *Xyletinus (Xyletinus) sveci* sp. nov. Other 25 species are noted as new to Greece - *Anobium hederæ* Ihsens, 1949; *A. inexpectatum* Lohse, 1954; *A. punctatum* (DeGeer, 1774); *Cacotemnus rufipes* (Fabricius, 1792); *Caenocara affine* (Sturm, 1837); *Dignomus jacqueti* (Pic, 1895); *Dorcatoma (Pilosodorcatoma) chrysolina* Sturm, 1837; *Episernus granulatus* Weise, 1887; *Ernobius lenkæ* Zahradník, 1996; *E. longicollis* (Sturm, 1837); *E. reitteri* Pic, 1902; *Gastrallus knizeki* Zahradník, 1996; *G. mauritanicus* Español, 1963; *G. vavrai* Zahradník, 2007; *Hadrobregmus denticollis* (Creutzer in Panzer, 1796); *Lasioderma bubalus phelipæarum* Peyerimhoff, 1926; *Ochina (Ochina) hirsuta* Seidlitz, 1889; *Oligomerus brunneus* (Olivier, 1790); *Priartobium serrifunus* Reitter, 1901; *Pseudoptilinus fissicollis* (Reitter in Saulcy, Reitter et Weise, 1877); *Ptinomorphus tatjanae* Logvinovskiy, 1978; *Ptinus (Pseudoptinus) nikitans* (Brenske et Reitter, 1884); *Stagetus byrrhoides* (Mulsant et Rey, 1861); *Xyletinus (Xyletinus) cylindricus* Kofler, 1970; *X. (X.) longitarsis* Jansson, 1942. The presence of the following five species in Greece is regarded as doubtful - *Dignomus aureopilis* (Desbrochers des Loges, 1875); *D. lesinae* (Reitter in Brenske et Reitter, 1884); *Eurostodes quisquilarium* (Baudi di Selve, 1874); *Ptinus (Ptinus) spitzyi spitzyi* A. Villa et G. B. Villa, 1838; *Pseudodryophilus paradoxus* (Rosenhauer, 1856). An annotated list of all 184 species known or mentioned from the area is given. Thirty-five species are endemic to Greece.

### INTRODUCTION

Despite numerous publications, Ptinidae of Greece have still been insufficiently known. Many species, including synonymized as well, were described from Greece (Boieldieu, 1854; Brenske et Reitter 1884; Fairmaire 1880; Gottwald 1977; Johnson 1975; Leiler 1979; Mařan 1941; Obenberger 1917; Pic 1896, 1901, 1902, 1903, 1934, Reitter 1884b, 1888, 1901; Roubal 1917a; Schilsky 1898, 1899, 1900; Toskina 2006, 2007a,b, 2011, 2012, 2013; Zahradník 1996; Zoufal 1897). The other species from Greece were presented in different catalogues (Fauvel 1889, Gemminger & Harold 1869; Heyden, Reitter & Weise 1883, 1891, 1906; Marseul 1886; Pic, 1912b; Jakobson 1913) or in publications addressed to single genera or genera groups (eg. Bellés 1996; Borowski 2002; Emets 1976; Emets & Logvinovskiy 1977; Español 1963a,b, 1964, 1967, 1969a,b, 1972, 1979a, 1992; Gottwald 1977; lablokoff-Khnzorian & Karapetyan 1991; Johnson 1975 etc.). Many data also come from papers dedicated to applied entomology (eg. Buchelos 1980, 1985, 1989; Buchelos & Athanassiou 1998; Howe 1958; Levinson & Buchelos 1988; Mourikis & Vassilaina-Alexopoulou 1975 etc.). The only overall work was the survey of genera of Greece and Crete (Oertzen 1886). Current survey of species living in Greece was provided by Borowski (2007) for subfamilies Gibbiinae a Ptinninae, and Zahradník (2007a) for the other subfamilies. There were mentioned 24 species in this work, but without further details.


At the present paper, a review of Greece Ptinidae is given. The present work is based on a literary search of almost 150 papers accompanied by study of nearly 500 specimens collected in Greece mainly by Czech, but also by Hungarian, German, Polish, Swiss and Swedish

entomologists, mainly over the last 25 years. The material contains 30 species new to Greece, including 5 specimens new to science described from Greece. Currently 184 specimens have been known from Greece, from which 35 specimens are endemic. Another 5 specimens were stated from Greece before based on wrong data and do not occur in Greece. The list of species as well as their descriptions are given below.

#### MATERIAL AND METHODS

All references given in the list under each species refer only to data from Greece. General distribution of all species in this work is given especially according to Borowski (2007) and Zahradník (2007a) and other sources from many papers. Locality data of other than type material are given usually in abbreviated form (Greece, province, prefecture, place - specific and extension, mount or mountain range, date of collecting, number of specimens, collector and holder of specimen/specimens), in alphabetical order. The abbreviation „spec.“ for specimen or specimens is used. The higher nomenclature (to genus or subgenus) used in this work is based on the publication Zahradník & Háva (2014), lower nomenclature (species, subspecies) is especially according to Borowski (2007) and Zahradník (2007a) and some other papers; all taxons are given in alphabetical order. Specimens from subfamily Ptininae were identified by J. Borowski (Rogow, Poland) and by author revid, other subfamilies were identified by the present author.

Map 1. Administrative division of Greece.  
(<http://www.recko-autem.cz/recko/zakladni-informace/uzemni-cleneni-recka>)



The specimens of the new species described are provided with a red, printed label showing the following text: „Holotype [Allotype or Paratype, respectively], on the second white label there is species name sp. n., P. Zahradník det.“. Description of new species contains in part of „Type material“ original writing of locality labels marked „xxx“; addition data are write in square brackets - [xxx]. All materials examined are marked by printed determination label with name of species, author and year of description, and P. Zahradník det. With hand-written year of description.

Acronyms of collection used in this paper are following:

HNHM Hungarian Natural History Museum, Budapest, Hungary;  
 NMP National Museum, Praha, Czech Republic;  
 USNM - Smithsonian Institution, Washington, D. C., USA.

## SPECIES ACCOUNT

### **ANOBIINAE FLEMING, 1821**

#### **Anobiini Fleming, 1821**

#### ***Anobium hederæ* Ihssens, 1949**

**Distribution.** Central and southern Europe.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Olympiada, vi.2008, 10 spec., ex larvae from *Hedera helix*, Z. Kraus leg., Z. Kraus et P. Zahradník coll. **New species for Greece.**

#### ***Anobium inexpectatum* Lohse, 1954**

**Distribution.** Central and southern Europe.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., ii.2011, 3 spec., J. Habarta leg., J. Habarta et P. Zahradník coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 12.-13.vi.2014, 1 spec., Z. Malinka leg. et coll.; Greece, Thessaly, Trikala pref., Kastraki, Kalambaka env., 1.vi.2012, 1 spec., Z. Malinka leg. et coll.

**A species new to Greece.**

#### ***Anobium punctatum* (DeGeer, 1774)**

**Distribution.** Cosmopolitan, originally European species.

**Material examined.** Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 4.-6.vi.2013, 5 spec., T. Sitek leg. et coll.

**New species for Greece.**

#### ***Cacotemnus rufipes* (Fabricius, 1792)**

**Distribution.** Europe, except southern area, Turkey, Siberia, Far East.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., iii. 2011, ex larvae, 1 spec., J. Habarta leg. et coll. **New species for Greece.**

#### ***Hemicoelus canaliculatus* (C. G. Thomson, 1863)**

*Hemicoelus canaliculatus* (C.G.Thomson, 1863): Zahradník, 2007a: 339.

*Anobium nitidum* Herbst, 1793: Reitter, 1901: 21; Jakobson, 1913: 816; Horion, 1961: 223.

*Byrrhus nitidus* (Herbst, 1793): Oertzen, 1886: 254.

**Distribution.** Europe, Far East, Iran.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.

***Hemicoelus costatus* (Aragona, 1830)**

*Anobium costatum* Aragona, 1830: Horion, 1961: 225.  
*Hemicoelus costatus* (Aragona, 1830): Zahradník, 2007a: 340.

**Distribution.** Europe, Turkey.

**Material examined.** Greece, Thessaly, Karditsa pref., Kiatra Broastra pass, Amarandos, 3.vi.2012, 2 spec., Z. Malinka leg. et coll.; Greece, Thessaly, Larisa pref., Spilia env., Ossa Oros Mt., 14.-17.vi.2010, 1 spec., J. Plecháč leg. et coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 29.-31.v.2012, 2 spec., T. Sitek leg. et coll.

***Hemicoelus fulvicornis* (Sturm, 1837)**

*Anobium fulvicorne* Sturm, 1837: Yakobson, 1913: 816; Horion, 1961: 224.  
*Hemicoelus fulvicornis* (Sturm, 1837): Zahradník, 2007a: 340.  
*Byrrhus fulvicornis* (Sturm, 1837): Oertzen, 1886: 254.

**Distribution.** Europe, Iran, Turkey.

**Material examined.** Greece, Central Macedonia, Imathia pref., Naoussa env., Vermio Mt., 6.-7.vi.2012, 1 spec., T. Sitek leg. et coll.; Greece, Central Macedonia, Serres pref., Ano Poroia, Kerkini Mt., 13.vi.2013, 2 spec., T. Sitek leg. et coll.; Greece, Eastern Macedonia and Thrace, Drama pref., Livadero, 31.v.2006, 4 spec., ex pupae from *Corylus avelana*, J. Ch. Vávra leg. et coll.; Greece, Thessaly, Larisa pref., Slomi, Ossa Oros Mt., 29.-31.v.2012, 3 spec., T. Sitek leg. et coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 30.v.2012, 1 spec., T. Sitek leg., P. Zahradník coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 4.-6.vi.2013, 3 spec., T. Sitek leg. et coll.

***Hemicoelus rufipennis* (Duftschmid, 1825)**

*Hemicoelus rufipennis* (Duftschmid, 1825): Zahradník, 2007a: 340.

**Distribution.** Central and southeastern Europe, Turkey.

**Material examined.** Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 4.-6.vi.2013, 1 spec., T. Sitek leg. et coll.

**Gastrallini White, 1982*****Falsogastrallus unistriatus* (Zoufal, 1897)**

*Gastrallus unistriatus* Zoufal, 1897: Zoufal, 1897: 206; Schilsky, 1898: 53; Schilsky, 1899: 36; Heyden, Reitter & Weise, 1906: 428; Pic, 1912b: 30; Yakobson, 1913: 818; Schaufuss, 1916: 722; Horion, 1972: 29; Logvinovskiy, 1985: 97.  
*Falsogastrallus unistriatus* (Zoufal, 1897): Español, 1970: 41; Halperin & Español, 1978: 10; Español, 1994: 107; Zahradník et al., 2000: 313; Toskina, 2003: 197; Zahradník, 2007a: 340; Zahradník, 2010: 272.

**Distribution.** Austria, Croatia, Greece, Hungary, Italy, Lebanon, Serbia, Syria, Tajikistan, Turkey.

**Material examined.** Greece, Western Greece, Achaea pref., Kalavryta, iv.1936, 4 spec., J. Mařan et D. Táborský leg., NMP coll.

***Gastrallus corsicus* Schilsky, 1898**

*Gastrallus corsicus* Schilsky, 1898: Zahradník, 2007a: 340.

**Distribution.** the Mediterranean.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.-20.vii.2002, 1 spec., Z. Kraus leg. et coll.; Greece, Crete, Chania pref., Chania, Agia Marina env., 15.-24.vi.2003, 1 spec., L. Daněk leg. et coll.; Greece, Crete, Lasithi pref., Mirtos, 1992, ex larvae, 1 spec., I. Jeniš leg., P. Zahradník coll.; Greece, Crete, Rethymno pref., Panormos, 12.vi.2004, 13 spec., 29.vi.2005, 14 spec., Z. Švec leg. et coll.; Greece, Crete, Rethymno pref., Rethymno, Adelianos Kampos, 7.-9.vi.2004, 3 spec., Z. Švec leg. et coll.; Greece, Crete, Rethymno pref., Scaleta, 16.vi.2004, 1 spec., Z. Švec leg. et coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Kolymbia, 23.-28.vi.2014, 1 spec., T. Staněk leg. et coll.; Greece, Thessaly, Larisa pref., Melivia, Ossa Oros Mt., 1.-4.vi.2007, 1 spec., G. Jarosiewicz leg., T. Gazurek coll.

***Gastrallus immarginatus* (P.W.J. Müller, 1821)**

*Gastrallus immarginatus* (P.W.J. Müller, 1821): Oertzen, 1886: 254; Yakobson, 1913: 818; Horion, 1961: 216; Logvinovskiy, 1985: 95; Zahradník, 2007a: 340.

**Distribution.** Europe, northern Africa, China.

***Gastrallus knizeki* Zahradník, 1996**

**Distribution.** Austria, Czech Republic, France, Germany, Slovakia.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 3 spec., J. Habarta leg. et coll.; Greece, Thessaly, Trikala pref., Kria Vrasi, v.2006, 6 spec., ex larvae from *Viscum album*, Z. Kraus leg., Z. Kraus et P. Zahradník coll. **New species for Greece.**

***Gastrallus kocheri* Español, 1963**

*Gastrallus kocheri* Español, 1963: Zahradník, 2007a: 340; Zahradník, 2007b: 173.

**Distribution.** Greece (Crete), Italy, Morocco, Spain, Tunisia.

**Material examined.** Greece, Crete, Lasithi pref., Myrtos, 3.vi.1990, 1 spec., I. Jeniš leg., P. Zahradník coll.

***Gastrallus laevigatus* (Olivier, 1790)**

*Gastrallus laevigatus* (Olivier, 1790): Horion, 1961: 218; Español, 1963a: 15; Zahradník, 2007a: 340.

**Distribution.** Europe, northern Africa, Middle East, Japan.

**Material examined.** Greece, Ionian Islands, Korfu pref., Korfu I., NMP coll.; Greece, Thessaly, Trikala pref., Kastraki, Kalambaka env., 11.-12.6.2010, 2 spec., T. Sitek leg. et coll.

***Gastrallus mauritanicus* Español, 1963**

**Distribution.** Italy, Morocco, Tunisia, Turkey.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Vrasna, vii.2004, 5 spec., viii.2007, 1 spec., all ex larvae from *Prunica* sp., Z. Kraus leg., Z. Kraus et P. Zahradník coll.; Greece, Crete, Rethymno pref., Bali, 17.-19.vii.2007, 2 spec., T. Růžička leg., T. Růžička et P. Zahradník coll. **New species for Greece.**

***Gastrallus vavrai* Zahradník, 2007**

**Distribution.** Croatia, Greece, Turkey.

**Material examined.** Greece, Crete, Lasithi pref., Mesavouni, Mt. Dikti, 10.vii.2007, 2 spec., D. Vacula leg., T. Sitek coll.; Greece, Thessaly, Trikala pref., Pinios riv., Kalambaka, 10.-11.vi.2010, 1 spec., Z. Malinka leg., P. Zahradník coll.; Greece, Western Greece, Aetolia-Akarnania pref., Náfpaktos env., 19.vi.2005, S. Benedikt leg. P. Zahradník coll. **New species for Greece.**

**Hadrobregmini White, 1982*****Hadrobregmus denticollis* (Creutzer in Panzer, 1796)**

**Distribution.** Europe (except eastern part).

**Material examined.** Greece, Central Greece, Fokida pref., Gravia env., 10.vi.2008, 1 spec., J. Mertlík leg., M. Zýka coll.; Greece, Eastern Macedonia and Thrace, Kavala pref., Akrovouni env., Pangeos Mt., 3.-4.vi.2013, 2 spec., T. Sitek leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kria Vrasi, Pindos Mt., 8.vi.2013, 1 spec., T. Sitek leg. et coll.

**New species for Greece.**

***Hadrobregmus pertinax* (Linnaeus, 1758)**

*Hadrobregmus pertinax* (Linnaeus, 1758): Zahradník, 2007a: 341.

**Distribution.** Europe, Far East, northern China.

**Material examined.** Greece, Epirus, Ioannina pref., Leptokaria, 30.vii.-15.viii.1993, 1 spec., J. Háva leg., P. Zahradník coll.

**Nicobiini White, 1982*****Nicobium castaneum* (Olivier, 1790)**

*Nicobium castaneum* (Olivier, 1790): Pic, 1912b: 32; Jakobson, 1913: 815; Horion, 1961: 219; Zahradník, 2007a: 342; Denux & Zagatti, 2010: 344.

*Anobium hirtum* Illiger, 1807: Schilsky, 1898: 31.

**Distribution.** Europe, Caucasus, northern Africa, Azores Islands, Canary Islands, Madeira Islands, China, Turkey.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodes I., 1 spec., NMP coll.

**Stegobiini White, 1982*****Oligomerus brunneus* (Olivier, 1790)**

**Distribution.** Europe, Siberia, Japan, Tadzhikistan.

**Material examined.** Greece, Thessaly, Thessaly, Trikala pref., Ambelia, Klinovitiko riv., Pindos Mt., 14.vi.2010, 1 spec., T. Sitek leg. et coll. **New species for Greece.**

***Oligomerus disruptus* (Baudi di Selve, 1874)**

*Oligomerus disruptus* (Baudi di Selve, 1874): Jakobson, 1913: 815; Zahradník, 2007a: 342.

*Byrrhus disruptus* (Baudi di Selve, 1874): Oertzen, 1886: 254;

**Distribution.** Algeria, Corsica Island, Cyprus, Greece, Italy.

***Oligomerus ptilinoides* (Wollaston, 1854)**

*Oligomerus ptilinoides* (Wollaston, 1854): Jakobson, 1913: 815; Horion, 1961: 213; Dominik, 1966: 34; Logvinovskiy, 1985: 92; Zahradník, 2007a: 342.

*Oligomerus reyi* (Brisout de Barneville in Grenier, 1867): Schilsky, 1898: 33.

*Byrrhus reyi* (Brisout de Barneville in Grenier, 1867): Oertzen, 1886: 254.

**Distribution.** Europe, northern Africa, Siberia, China, Japan, Syria, Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Central Macedonia, Pieria pref., Litochoro, 8.viii.1997, 1 spec., M. Mantič leg. et coll.; Greece, Crete, Chania pref., Hania env., Omalos plain, ix.2003, 1 spec., ex larvae, J. Batelka & H. Batelková leg., J. Batelka coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Rhodos env., vi.2009, A. Kotána leg., HNHM coll.

***Oligomerus retowskii* Schilsky, 1898**

*Oligomerus retowskii* Schilsky, 1898: Schilsky, 1898: 32; Reitter, 1901: 22; Heyden, Reitter & Weise, 1906: 429; Pic, 1912b: 27; Jakobson, 1913: 815; Schaufuss, 1916: 725; Zaytsev, 1956: 58; Español, 1981: 49; Logvinovskiy, 1985: 92; Zahradník, 1998: 271; Zahradník, 2007a: 342.

**Distribution.** Caucasus, southern Balkan Peninsula, Ukraine, southern Russia.

**Priartobium serrifunis Reitter, 1901**

**Distribution.** Algeria, Bulgaria, Corse Island, Croatia, Israel, Italy, Ukraine.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., Kolymbia, 22.-28.vi.2014, 1 spec., T. Staněk leg. et coll. **New species for Greece.**

**Stegobium paniceum (Linnaeus, 1758)**

*Stegobium paniceum* (Linnaeus, 1758): Yakobson, 1913: 815; Buchelos, 1985: 222; Buchelos & Anthanassiou, 1998: 131; Zahradník, 2007a: 343.

*Anobium paniceum* (Linnaeus, 1758): Rey, 1892: 2.

*Sitodrepa panicea* (Linnaeus, 1758): Sahlberg, 1903b: 64; Pelekassis, 1962: 67.

*Byrrhus paniceus* (Linnaeus, 1758): Oertzen, 1886: 254.

*Anobium minutum* Fabricius, 1792: Kiesenwetter, 1859: 162.

**Distribution.** Cosmopolitan, originally most likely to come from Mediterranean area.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Meteora, 24.-30.v.2007, 1 spec., Dolinská leg., E. Ezer coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.-20.vii.2002, 1 spec., Z. Kraus leg. et coll.; Greece, Crete, Rethymno pref., Plakias env., 6.-16.vii.2005, 1 spec., P. Zahradník leg et coll.; Greece, Ionian Islands, Kefalonia and Ithaca prov., Kefalonia I., Poros env., 30.v.-5.vi.1997, 1 spec., K. Schön leg., P. Zahradník coll.

**DORCATOMINAE C.G. THOMSON, 1859****Dorcatomini C.G. Thomson, 1859****Anitys rubens (Hoffmann, 1803)**

*Anitys rubens* (Hoffmann, 1803): Logvinovskiy, 1985: 159; Zahradník, 2007a: 343.

**Distribution.** European (except western Mediterranean area), Egypt.

**Caenocara affine (Sturm, 1837)**

**Distribution.** Europe, Far East.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kastraki, 1.vi.2012, 1 spec., Z. Malinka leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kria Vrasi Pindos Mt., 8.vi.2013, 1 spec., T. Sitek leg. et coll. **New species for Greece.**

**Caenocara subglobosum (Mulsant et Rey, 1864)**

*Caenocara subglobosum* (Mulsant et Rey, 1864): Oertzen, 1886: 255; Yakobson, 1913: 811; Horion, 1961: 253; Logvinovskiy, 1985: 160; Zahradník, 2007a: 344.

**Distribution.** Central and northern Europe, Caucasus, northern Africa, Siberia, Far East.

**Dorcatoma (Dorcatoma) palmi Zahradník, 1996**

*Dorcatoma (Dorcatoma) palmi* Zahradník, 1996: Zahradník, 1996: 98; Zahradník, 2007a: 344.

**Distribution.** Cyprus, Greece.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., 24.vi.-4.vii.1958, 6 spec. (paratypes), T. Palm leg., P. Zahradník et USNM coll.

**Dorcatoma (Pilodorcatoma) chrysomelina Sturm, 1837**

**Distribution.** Central and northern Europe, Siberia, Far East.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Vrasna, 27.v.2007, 1 spec., Z. Kraus leg., P. Zahradník coll.; Greece, Thessaly, Larisa pref., Olympiada, 11.-30.vii.2004, 1 spec., Z. Kraus leg. et coll.; Greece, Thessaly, Larisa pref., Stomio env., Ossa Oros Mt., 13.vi.2014, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Trikala pref., Ambelia, Klinovitiko riv., Pindos Mt., 14.vi.2010, 1 spec., T. Sitek leg., P. Zahradník coll. **New species for Greece.**

### ***Dorcatoma (Pilosodorcatoma) setosella setosella* Mulsant et Rey, 1864**

*Dorcatoma (Pilosodorcatoma) setosella setosella* Mulsant et Rey, 1864: Oertzen, 1886: 255; Schilsky, 1899: 62; Pic, 1912b: 75; Yakobson, 1913: 811; Horion, 1961: 252; Logvinovskiy, 1980: 152; Zahradník, 2007a: 344.

**Distribution.** Central and western Europe, Greece, Turkey.

**Material examined.** Greece, Western Greece, Achaia pref., Kalavryta, vi.1997, ex larvae from *Phelinus* sp. to *Prunus*, 1 spec., B. Büche leg. et coll.

## **Prothecini White, 1982**

### ***Stagetus andalusiacus cribricollis* (Aubé, 1861)**

*Stagetus andalusiacus cribricollis* (Aubé, 1861): Zahradník, 2007a: 346.  
*Theca remota* Brenske et Reitter, 1884: Pic, 1912b: 64.

**Distribution.** Northern Africa, Greece, Italy.

### ***Stagetus byrrhoides* (Mulsant et Rey, 1861)**

**Distribution.** Central and southern Europe, northern Africa, Middle East.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., 24.vi.-4.vii.1958, 1 spec., T. Palm leg., P. Zahradník coll. **New species for Greece.**

### ***Stagetus conicollis* (Schilsky, 1899)**

*Stagetus conicollis* (Schilsky, 1899): Yakobson, 1913: 810; Schaufuss, 1916: 727; Español, 1969b: 57; Zahradník, 2007a: 346.  
*Theca conicollis* Schilsky, 1899: Schilsky, 1899: 48; Reitter, 1901: 41; Pic, 1912b: 62.

**Distribution.** Greece.

### ***Stagetus curimoides curimoides* (Reitter in Brenske et Reitter, 1884)**

*Stagetus curimoides curimoides* (Reitter in Brenske et Reitter, 1884): Heyden, Reitter & Weise, 1906: 432; Sahlberg, 1913: 17; Yakobson, 1913: 810; Schaufuss, 1916: 727; Chikatunov, 2000: 61; Zahradník et al., 2000: 316; Zahradník, 2007a: 346.

*Theca curimoides* Reitter in Brenske et Reitter, 1884: Brenske et Reitter, 1884: 86; Marseul, 1886: 298; Oertzen, 1886: 255; Heyden, Reitter & Weise, 1891: 233; Schilsky, 1899: 45; Reitter, 1901: 40; Sahlberg, 1903b: 64; Pic, 1912b: 62.

**Distribution.** Middle East, Greece.

**Material examined.** Greece, Ionian Islands, Zakynthos pref., Zakynthos I., Katastari, 23.iv.1971, 1 spec., B. Hauser leg., P. Zahradník coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Kattavia, 12.iv.1977, 1 spec., J.-C. Besuchet leg., P. Zahradník coll.

### ***Stagetus curimoides exiguus* Roubal, 1917**

*Stagetus curimoides exiguus* Roubal, 1917: Roubal, 1917a: 53; Zahradník, 2007a: 346.



**Distribution.** Greece.

### ***Stagetus elongatus* (Mulsant et Rey, 1861)**

*Stagetus elongatus* (Mulsant et Rey, 1861): Español, 1969a: 115; Zahradník, 2007a: 346.

*Stagetus rufescens* (Pic, 1902): Heyden, Reitter & Weise, 1906: 432; Yakobson, 1913: 810; Schaufuss, 1916: 727.

*Theca rufescens* Pic, 1902: Pic, 1902a: 80.

**Distribution.** Southern and southwestern Europe, northern Africa, Middle East, Turkey.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.-20.vii.2002, 1 spec., Z. Kraus leg. et coll.; Greece, Crete, Heraklio pref., Mesavouni, Mt. Dikti, 10.vii.2007, 2 spec., D. Vacula leg., T. Sitek coll.; Greece, Epirus, Preveza pref., Preveza env., 6.-16.vi.1992, 1 spec., T. Růžička leg., P. Zahradník coll.; Greece, Ionian Islands, Corfu pref., Corfu I., Benitses env., 23.-29.vi.2003, 1 spec., K. Orszulik leg. et coll.; Greece, Thessaly, Trikala pref., Ambelia, Klinovitiko riv., Pindos Mt., 14.vi.2010, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka, Pinios riv., 10.-11.vi.2010, 1 spec., T. Sitek leg.; P. Zahradník coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kastraki, 11.-12.vi.2010, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Karditsa pref., Rentina, Modi, 3.-20.vii.2002, 1 spec., Z. Kraus leg. et coll.; Greece, Western Macedonia, Grevena pref., Karpero env., Anixi, 10.vi.2013, 10 spec., T. Sitek leg. et coll.; Greece, Western Greece, Achaia pref., Patra env., Metohi, 15.vi.2005, 1 spec., J. Plecháč leg., P. Zahradník coll.; Greece, Western Macedonia, Grevena pref., Karpero env., Anixi, 18.vi.2014, 4 spec., T. Sitek leg. et coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, vii.2004, 1 spec., ex larvae from *Prunica* sp., Z. Kraus leg. et coll.

### ***Stagetus espanoli* Toskina, 2012**

*Stagetus espanoli* Toskina, 2012: Toskina, 2012: 19.

**Distribution.** Greece.

### ***Stagetus franzi* Español, 1969**

*Stagetus franzi* Español, 1969: Español, 1969a: 117; Español, 1969b: 57; Español, 1977: 10; Zahradník, 2007a: 346.

**Distribution.** Northern Africa, Middle East, France, Greece, Hungary, Spain, Turkey.

**Material examined.** Greece, Peloponnese, Lakonia pref., Kelefa, 11.v.2009, 1 spec., O. Konvička leg. et coll.

### ***Stagetus italicus* (Reitter, 1885)**

*Stagetus italicus* (Reitter, 1885): Heyden, Reitter & Weise, 1906: 432; Yakobson, 1913: 810; Schaufuss, 1916: 727; Zahradník, 2007a: 346.

*Theca italica* Reitter, 1885: Heyden, Reitter & Weise, 1891: 233; Reitter, 1901: 42; Pic, 1912b: 63.

**Distribution.** Greece, Italy.

### ***Stagetus pilula* (Aubé, 1861)**

*Stagetus pilula* (Aubé, 1861): Schilsky, 1899: 50; Heyden, Reitter & Weise, 1906: 432; Yakobson, 1913: 810; Schaufuss, 1916: 727; Horion, 1961: 245; Logvinovskiy, 1985: 149; Zahradník, 2007a: 347.

*Theca pilula* Aubé, 1861: Oertzen, 1886: 255; Reitter, 1901: 41; Sahlberg, 1903b: 64; Pic, 1912b: 63.

**Distribution.** Central and southern Europe, northern Africa, Middle East, Turkey.

### ***Stagetus propinquus* Obenberger, 1917**

*Stagetus propinquus* Obenberger, 1917: Obenberger, 1917: 36; Zahradník, 2007a: 347.

**Distribution.** Greece.

***Stagetus puncticollis* (Brenske et Reitter, 1884)**

*Stagetus puncticollis* (Brenske et Reitter, 1884): Heyden, Reitter & Weise, 1906: 432; Yakobson, 1913: 810; Schaufuss, 1916: 727; Español, 1969a: 113; Zahradník, 2007a: 347.

*Theca puncticollis* Brenske et Reitter, 1884: Reitter, 1884a: 120; Oertzen, 1886: 255; Heyden, Reitter & Weise, 1891: 233; Schilsky, 1899: 49; Pic, 1912b: 63.

*Theca moreana* Pic, 1921: Pic, 1921b: 9.

**Distribution.** Algeria, Cyprus, Greece, Israel, Morocco, Portugal, Tunisia, Turkey.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., Lardos, 1.vii.2007, 1 spec., K. Orszulik leg. et coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Paradissi env., Paradissi Mt., 5.v.1996, 1 spec., K.Schön leg., P. Zahradník coll.

**DRYOPHILINAE GISTL, 1848****Dryophilini Gistl, 1848*****Dryophilus densipilis* Abeille de Perrin, 1872**

*Dryophilus densipilis* Abeille de Perrin, 1872: Zahradník, 2007a: 348.

**Distribution.** Southern Europe, Algeria.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., Ialysos, Filempos Mt., 10.v.1996, 1 spec., R. Borovec leg., J. Ch. Vávra coll.

***Dryophilus longicollis* (Mulsant et Rey, 1853)**

*Dryophilus longicollis* (Mulsant et Rey, 1853): Zahradník, 2007a: 348.

*Dryophilus longicollis elongatus* Pic, 1921: Pic, 1921a: 2.

**Distribution.** Central, southern and western Europe, northern Africa.

***Dryophilus pusillus* (Gyllenhal, 1808)**

*Dryophilus pusillus* (Gyllenhal, 1808): Oertzen, 1886: 254; Yakobson, 1913: 818; Logvinovskiy, 1985: 48; Zahradník, 2007a: 348.

*Anobium pusillum* Gyllenhal, 1808: Kiesenwetter, 1859: 162.

**Distribution.** Europe, Caucasus, Egypt.

***Dryophilus rufescens* Pic, 1921**

*Dryophilus rufescens* Pic, 1921: Pic, 1921a: 2; Zahradník, 2007a: 348.

**Distribution.** Greece.

**Material examined.** Greece, Crete, Heraklio pref., Thomadiano, Viannos, 7.-8.iv.2004, 1 spec., J. Batelka & H. Batelková leg., P. Zahradník coll.; Greece, Northern Aegean islands, Lesbos pref., Ano Chorio env., Thripti Mt., 23.iii.2014, 1 spec., S. Benedikt leg., V. Benedikt coll.; Greece, Northern Aegean islands, Lesbos pref., Kallonii, 22.iv.2007, 1 spec., T. Růžička leg., P. Zahradník coll.

***Homophthalmus rugicollis* (Mulsant et Rey, 1853)**

*Dryophilus rugicollis* (Mulsant et Rey, 1853): Oertzen, 1886: 254; Yakobson, 1913: 818; Horion, 1961: 195; Novak, 1964: 87; Logvinovskiy, 1985: 46; Burakowski, Mroczkowski & Stefańska, 1986: 52.

*Homophthalmus rugicollis* (Mulsant et Rey, 1853): Zahradník, 2007a: 348.

*Dryophilus pusilloides* Obenberger, 1917: Obenberger, 1917: 35.

**Distribution.** Southern Europe, Caucasus, Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Central Macedonia, Thessaloniki pref., Stavros, 8.-10.vi.2014, 1 spec., Z. Malinka leg. et coll.; Greece, South Aegean islands, Naxos pref., Galini, 25.iv.1997, 1 spec., Lienland leg., P. Zahradnik coll.; Greece, Thessaly, Karditsa pref., Neraida, Plastiras lake, 1.vi.2013, 1 spec., O. Konvička leg. et coll.

### ***Pseudodryophilus paradoxus* (Rosenhauer, 1856)**

*Dryophilus paradoxus* (Rosenhauer, 1856): Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 230; Heyden, Reitter & Weise, 1906: 427; Pic, 1912b: 11; Jakobson, 1913: 818; Schaufuss, 1916: 722; Logvinovskiy, 1985: 46.

*Pseudodryophilus paradoxus* (Rosenhauer, 1856): Español, 1992: 42 (?).

**Distribution.** Western Mediterranean area.

**Note:** According to Zahradnik (2007) this species does not occur in Greece.

## **ERNOBIINAE PIC, 1912**

### **Ernobiini Pic, 1912**

#### ***Episernus graecus* sp. nov.**

(Figs. 1-3)

**Type material.** Holotype (male): „Greece - Peloponnese, Erimantos mts., Kalendzi env. [Western Greece, Achaia pref.], [poklep suché jedle] [knocking down from died fire], 15.5.2007, K. Rébl lgt.“, P. Zahradnik coll.

**Description.** Male (holotype). Elongate-oval, transversally slightly convex. Body length 2.6 mm, the greatest width 1.0 mm. Ratio elytra length : elytra width of 2.0. Piceous black. Antennae, palpi and legs yellow-brown. Pubescence white.

Head convex, shining, coarsely and densely punctate, pubescence fine, short, sparse, inclined forwards. Eyes strongly convex, globular, glabrous. Front 2.9 times wider than width of eye in dorsal view. Antennae consisting of ten antennomeres, filiform (Fig. 1). The first antennomere robust, 3 times as long as wide, the second 1.2 times longer than wide, the same width as previous. The third antennomere narrower, of the same length as the second. The fourth slightly shorter than the third, of the same width. The fifth and the sixth as long as wide, the seventh slightly transverse, shorter than previous. The last three antennomeres enlarged. The eighth of the same length as the third to the seventh together. The ninth slightly longer than eighth, the tenth 1.3 times longer than ninth. The last maxillary palpomere sharpened.

Pronotum transverse, ratio length : width 0.8, the greatest width shortly before base. Pronotum in middle with „V“ shaped flattened bump, and on slant edge in posterior angles (Fig. 2). Surface of pronotum shining, densely and coarsely punctate, distance between punctures the same as their diameter. Pubescence fine, short, sparse, inclined backwards. Scutellum small, square.

Elytra shortly oval, without distinct shoulders, shining, densely and coarsely punctate; distance between punctures the same as their diameter. Punctures not arranged in regular rows. Pubescence sparse, very short, recumbent, inclined backward.


Legs thin and long. Tibiae as long as tarsi. The first tarsomere the longest, 1.4 times longer than the second. The third and the fourth of the same length, together of the same length as the second. The fifth twice longer than the fourth. All tarsomere of the same width. Claws long and slim. The first ventrite in middle very narrow, on the sides twice longer than in middle. Other ventrites of the same length, slightly longer than the first on lateral margin.

Aedeagus see Fig. 3.

Female. Unknown.

**Differential diagnosis.** From other species of the genus *Episernus* differs by „V” shaped flattened bump on pronotum and by shape of aedeagus.

**Name derivation.** Derived from the name of country where the species was collected.



Figs. 1-3. *Episernus graecus* sp. nov.: 1- antenna of male; 2- pronotum; 3- aedeagus.

### ***Episernus granulatus* Weise, 1887**

**Distribution.** Central and western Europe, Siberia.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Fourka env., Smolikas Mt., 5.vi.2008, 2 spec., K. Rébl leg., K. Rébl et P. Zahradník coll. **New species for Greece.**

### ***Episernus taygetanus* Mařan, 1941**

*Episernus taygetanus* Mařan, 1941: Mařan, 1941: 123; Zahradník, 2007a: 349.

**Distribution.** Greece.

### ***Ernobius abietinus* (Gyllenhal, 1808)**

*Ernobius abietinus* (Gyllenhal, 1808): Kiesenwetter, 1877: 121; Oertzen, 1886: 254; Yakobson, 1913: 817; Horion, 1961: 207; Zahradník, 2007a: 349.

*Anobium abietinum* Gyllenhal, 1808: Kiesenwetter, 1859: 162.

**Distribution.** Europe (except bigger part of southern part), Caucasus.

### ***Ernobius abietis* (Fabricius, 1792)**

*Ernobius abietis* (Fabricius, 1792): Oertzen, 1886: 254; Horion, 1961: 208; Cymorek, 1974: 64; Johnson, 1975: 85; Burakowski, Mroczkovski & Stefańska, 1986: 56; Zahradník, 2007a: 349.

**Distribution.** Europe, Palearctic Asia, Algeria.

***Ernobius angusticollis* (Ratzeburg, 1837)**

*Ernobius angusticollis* (Ratzeburg, 1837): Logvinovskiy, 1977a: 126; Roques, 1983: 52; Zahradnik, 2007a: 349.

**Distribution.** Europe (except bigger area of southern part).

***Ernobius cupressi* Chobaut, 1899**

*Ernobius cupressi* Chobaut, 1899: Roques, 1983: 107; Zahradnik, 2007a: 349.

**Distribution.** Cyprus, France, Greece, Spain.

***Ernobius kailidisi* Johnson, 1975**

*Ernobius kailidisi* Johnson, 1975: Johnson, 1975: 86; Español, 1977: 8; Zahradnik, 1998: 268; Zahradnik, 2007a: 350.

**Distribution.** Greece, Turkey.

**Material examined.** Greece, Central Greece, Fthiotida pref., Aranova, Parnassus Mt., 6.-14.vi.1989, 1 spec., V. Němec leg., P. Zahradnik coll.; Greece, Central Greece, Fthiotida pref., Aranova, Parnassus Mt., 5.v.1994, 1 spec., I. Jeniš leg., P. Zahradnik coll.; Greece, Peloponnese, Lakonia pref., Knoneri, Taygetos Mt., 29.iv.2008, 1 spec., S. Benedikt leg., P. Zahradnik coll.; Greece, Thessaly, Larisa pref., Spifia, Ossa Oros Mt., 1.-2.vi.2007, 3 spec., T. Sitek leg., T. Sitek et P. Zahradnik coll.

***Ernobius kiesenwetteri* Schilsky, 1898**

*Ernobius kiesenwetteri* Schilsky, 1898: Johnson, 1975: 74; Zahradnik, 2007a: 350.

*Ernobius schilskyi* Reitter, 1901: Reitter, 1901: 15; Heyden, Reitter & Weise, 1906: 428; Pic, 1912b: 23; Yakobson, 1913: 816; Pic, 1916: 4; Hubenthal, 1927: 84; Cymorek, 1974: 62.

**Distribution.** Central and southern Europe, Caucasus, Turkey.

**Material examined.** Greece, Peloponnese, Arkadia pref., Menalo Mt., 1.-5.vi.2009, 9 spec., Z. Švec leg. et coll.; Greece, Western Greece, Achaia pref., Chelmos Mt., 23.v.2010, 3 spec., K. Rébl leg., K. Rébl et P. Zahradnik coll.

***Ernobius laticollis* Pic, 1927**

*Ernobius laticollis* Pic, 1927: Zahradnik, 2007a: 350.

**Distribution.** Austria, France, Greece, Italy, Spain, Switzerland, Tunisia.

**Material examined.** Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 1 spec., Z. Švec leg., P. Zahradnik coll.

***Ernobius lenkae* Zahradnik, 1996**

**Distribution.** Turkey.

**Material examined.** Greece, Crete, Heraklio pref., Pefkos 4.-5.vi.2004, 1 spec., V. Němec leg., P. Zahradnik coll.

**New species for Greece.**

***Ernobius longicollis* (Sturm, 1837)**

**Distribution.** Central, northern and western Europe, Siberia.

**Material examined.** Greece, Epirus, Ioannina pref., Metsovo, Katara pass, Indos Mt., 29.-30.v.2006, 1 spec., Z. Kraus leg. et coll. **New species for Greece.**

***Ernobius mollis mollis* (Linnaeus, 1758)**

*Ernobius mollis mollis* (Linnaeus, 1758): Oertzen, 1886: 254; Yakobson, 1913: 817; Sharp, 1916: 219; Markalas, 1991: 74; Zahradník, 2007a: 350.

**Distribution.** Europe, northern Africa, Siberia, Far East, China.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Fourka env., Smolikas Mt., 5.vi.2008, 4 spec., K. Rébl leg., K. Rébl et P. Zahradník coll.; Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 2 spec., Z. Švec leg. et coll.; Greece, Crete, Chania pref., Omolos, 22.-30.v.1990, 1 spec., I. Jeniš leg., P. Zahradník coll.; Greece, Crete, Lasithi pref., Anatoli env., 26.iii.2014, 1 spec., S. Benedikt leg., V. Benedikt coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Kolymbia, 23.-27.vi.2014, 1 spec., T. Staněk leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka, Kria Vrisi, Pindos Mt., 12.vi.2010, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Larisa pref., Spilia env., Ossa Oros Mt., 14.-17.vi.2010, 1 spec., J. Plecháč leg. et coll.; Greece, Western Greece, Elis pref., Vartholomio env., Glifa, 22.iv.2008, 1 spec., S. Benedikt leg.; P. Zahradník coll.

***Ernobius nigrinus* (Sturm, 1837)**

*Ernobius nigrinus* (Sturm, 1837): Zahradník, 2007a: 350.

**Distribution.** Europe, Canary Islands, Turkey.

**Material examined.** Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 1 spec., Z. Švec leg. et coll.; Greece, Central Macedonia, Pieria pref., Litochoro, Olympos Mt., 17.-18.vi.2010, 1 spec., J. Plecháč leg. et coll.

***Ernobius oertzeni* Schilsky, 1900**

*Ernobius oertzeni* Schilsky, 1900: Schilsky, 1900: 14; Reitter, 1901: 19; Heyden, Reitter & Weise, 1906: 429; Pic, 1912b: 22; Yakobson, 1913: 817; Schaufuss, 1916: 719; Cymorek, 1974: 62; Johnson, 1975: 91; Halperin & Español, 1978: 7; Roques, 1983: 96; Zahradník, 2000: 133; Zahradník et al., 2000: 312; Zahradník, 2007a: 351.

*Ernobius schilskyanus* Sharp, 1916: Sharp, 1916: 221.

**Distribution.** Cyprus, Greece, Israel, Italy.

***Ernobius pallidipennis* Pic, 1902**

*Ernobius pallidipennis* Pic, 1902: Zahradník, 2007a: 351.

**Distribution.** Algeria, Greece, Morocco, Spain.

**Material examined.** Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 7 spec., Z. Švec leg., Z. Švec et P. Zahradník coll.; Greece, Central Macedonia, Pieria pref., Litochoro env., Olympos Mt., 6 spec., J. Plecháč leg., P. Zahradník coll.

***Ernobius pini pini* (Sturm, 1837)**

*Ernobius pini* (Sturm, 1837): Johnson, 1975: 92; Logvinovskiy, 1977a: 126; Roques, 1983: 86; Zahradník, 2007a: 351.

**Distribution.** Europe, northern Africa, Turkey.

**Material examined.** Greece, Eastern Macedonia and Thrace, Kavala pref., Kalivia, Vathalomio, 22.v.2007, 1 spec., S. Benedikt leg., P. Zahradník coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Salakos env., Profitis Illias Mt., 6.v.1996, 1 spec., K. Schön leg., P. Zahradník coll.

***Ernobius pruinosus* (Mulsant et Rey, 1863)**

*Ernobius pruinosus* (Mulsant et Rey, 1863): Zahradník, 2007a: 351.

**Distribution.** Algeria, France, Germany, Greece, Italy, Lebanon, Morocco, Slovakia, Spain, Tunisia.

**Material examined.** Greece, Central Greece, Euboia pref., Stropones, Col de Streni, 30.iv.1987, 2 spec., Lienhard leg., P. Zahradník coll.; Greece, Central Greece, Evrytania pref., Aniada env., Kaliakouda Mt., 27.vi.2005, 1 spec., S. Benedikt leg., P. Zahradník coll.

***Ernobius rebli* sp. nov.**

(Figs. 4-7)

**Type material.** Holotype (male): „Greece - Peloponnese b., Metochi [Western Greece, Achaia pref.], 14.5.2007, K. Rébl lgt.“; P. Zahradník coll. Allotype (female): the same data as holotype; P. Zahradník coll. Paratypes: 1 male, the same data as holotype; K. Rébl coll.; 1 female: „Greece - Peloponnese, Erimantos mts., Kalendzi env., (poklep suché jedle) [knocking down of died fire], 15.5.2007, K. Rébl lgt.“, P. Zahradník coll.

**Description.** Male (holotype). Elongate-oval, transversally slightly convex. Body length 3.1 mm, the greatest width 1.2 mm. Ratio elytra length : elytra width of 1.9. Yellow - brown, including antennae, palpi and legs. Pubescence golden-yellow, recumbent.

Head evenly convex, shining, coarsely and densely punctate, with sparse recumbent pubescence, inclined forwards. Eyes strongly convex, globular, glabrous. Front 2.7 times wider than width of eye in dorsal view. Antennae consisting of eleven antennomeres, filiform (Fig. 4a). The first antennomere robust, twice longer than wide. The second shortly oval. The third and fourth narrower than second, shortly longer than wide, both almost of the same length, the fourth minutely shorter. The fifth (and other) slightly wider, twice longer than fourth. The sixth short, only one half of previous. The seventh of the same length as the fifth, but slightly wider. The eighth shorter, only 2/3 of previous and also slightly narrower. Antennomeres nine, ten and eleven enlarged. The ninth shorter than sixth to eighth antennomeres together, slightly shorter than the tenth or eleventh. The tenth of the same length as seventh and eighth together and of the same length as the eleventh. The last maxillary palpomeres sharpened.

Pronotum transverse (Fig 5.), ratio length : width 0.7, shining, disc slightly convex, lateral margin slightly and narrowly flattened. All pronotal angles rounded. Base of pronotum almost straight. Surface of pronotum coarsely and densely punctate, punctures circular. Pubescence recumbent, long, sparse, inclined backward, on sides of pronotum askew from central line of pronotum. Scutellum small, slightly transverse, trapezoidal, on base of pronotum slightly longer than on apex.

Elytra shortly oval, without distinct shoulders, shining, densely and coarsely punctate; punctures circular, distance between them the same as their diameter. Punctures not arranged in regular rows. Pubescence sparse, long, recumbent, inclined backward. Apex of each elytron reddish.

Legs thin and long. Tibiae as long as tarsi. The first tarsomere 1.2 times longer than the second. The third tarsomere half length as previous. The fourth deeply heart-shapedly emarginate, up to two thirds length of emargination, the fifth tarsomere is inserted, which is lengthily oval, three times longer as previous. Claws relatively large, without teeth (Fig. 6).

Disc of metasternum transversally slightly convex, posteriorly with small rounded depression. Surface shining, densely and coarsely umbilicate punctate, Puncture diameter as long as distance between punctures. Pubescence sparse, fine, short, recumbent, inclined backwards. Ventriles shining, densely and coarsely, umbilicate punctate; puncture diameter as long as distance between punctures. Pubescence sparse, fine, short, recumbent, inclined backwards, too.


Aedeagus see Fig. 7.

Female (allotype). Bigger, body length 3.7 mm, the greatest width 1.4 mm. Sexual dimorphism visible on antennae. Female has shorter last three antennomeres (ratio between them is the same), antennomere seventh and eight slightly longer the ninth. Antennomeres fifth and seventh of the same length as sixth and eighth (Fig. 4b).

**Variability.** Body length 3.0 - 3.2 mm, the greatest width 1.1 - 1.3 mm. Reddish apex sometimes invisible.

**Differential diagnosis.** This species belongs to *Ernobius pini*-group. *E. rebli* sp. nov. differs from *E. madoni*, *E. rufus*, and *E. pini* by presence of only circular punctures. Species *E. oertzeni*, *E. parens*, *E. gallicus*, and *E. fulvus*, have ninth antennomere as long as the seventh and the eighth together. From *E. impressithorax*, which have ninth antennomere longer than the seventh and the eighth (as *E. rebli*) it differs by sparser punctuation, elytra are more shining, and pronotum is not too flattened. The shape of aedeagus is quite different.

**Name derivation.** Dedicated to the collector of the type series, Karel Rébl, well-known entomologist and specialist in the genus Bembidion (Carabidae).



Figs. 4-7. *Ernobius rebli* sp. nov.: 4a- antenna of male; 4b- antenna of female, 5- pronotum; 6- back tarsus; 7- aedeagus.

### ***Ernobius reitteri* Pic, 1902**

**Distribution.** Turkey.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Rhodos I., Psintos, 30.ix.-5.x.2009, 2 spec., D. Vacula leg., P. Zahradník coll. **New species for Greece.**

### ***Ernobius robusticornis* Mařan, 1941**

*Ernobius robusticornis* Mařan, 1941: Mařan, 1941: 124; Johnson, 1975: 88; Zahradník, 1998: 267; Zahradník, 2007a: 351.

**Distribution.** Greece, Turkey.



**Ochinini Zahradnik in Zahradnik et Háva, 2014*****Ochina (Dulgieris) latreillii (Bonelli, 1812)***

*Ochina latreillii* (Bonelli, 1812): Schaufuss, 1916: 727; Horion, 1961: 198; Español, 1963b: 292; Jelínek & Rébl, 1990: 235; Cecchi & Nardi, 1998: 41; Zahradnik, 2007a: 352.

**Distribution.** Central and southeastern Europe, Caucasus, Turkey.

**Material examined.** Greece, Epirus, Ioannina pref., Metsovo env., Notia Pindos Mt., 13.vi.2012, 1 spec., B. Zbuzek leg. et coll.

***Ochina (Ochina) ferruginea Schilsky 1899***

*Ochina ferruginea* Schilsky 1899: Schilsky, 1899: 11; Reitter, 1901: 37; Heyden, Reitter & Weise, 1906: 432; Pic, 1908: 42; Pic, 1912b: 12; Yakobson, 1913: 812; Porta, 1929: 489; Español, 1963b: 287; Logvinovskiy, 1977a: 127; Karapetyan, 1984: 948; Logvinovskiy, 1985: 63; Zahradnik, 2007a: 352.

*Ochina (Ochina) pallidicolor* (Pic, 1908): Schaufuss, 1916: 727 (it is mistake; this species was described by Pis from Roma, Italy).

**Distribution.** Azerbaijan, Greece, Italy.

***Ochina (Ochina) hirsuta Seidlitz, 1889***

**Distribution.** Algeria, France, Italy, Morocco, Spain, Tunisia.

**Material examined.** Greece, Ionian Islands, Corfu pref., Lefkimmí env., 15.vi.2002, 1 spec., K. Schön leg., P. Zahradnik coll.; Greece, Ionian Islands, Corfu pref., Sidari, 27.v.1997, 2 spec., J. Hájek leg., NMP et P. Zahradnik coll. **New species for Greece.**

***Ochina (Ochina) ptinoides (Marsham, 1802)***

*Ochina ptinoides* (Marsham, 1802): Yakobson, 1913: 812; Horion, 1961: 199;

*Ochina hederæ* (P.W.J.Müller, 1821): Oertzen, 1886: 254; Zahradnik, 2007a: 353.

**Distribution.** Europe, northern Africa, Turkey.

**Material examined.** Greece, Crete, Rethymno pref., Platania, 23.v.2006, 1 spec., M. Šárovec leg., P. Zahradnik coll.

**Xestobiini Böving, 1927*****Hyperisus plumbeum (Illiger, 1801)***

*Hyperisus plumbeum* (Illiger, 1801): Zahradnik, 2007a: 352.

*Xestobium plumbeum* (Illiger, 1801): Español, 1964: 125.

*Xestobium graecum* Mařan, 1941: Mařan, 1941: 122.

**Distribution.** Central and southern Europe, Caucasus, Syria, Turkey.

**Material examined.** Greece, Crete, Chania pref., Omalos, Lefka Ori 15.v.2010, 3 spec., ex larvae from Zelkova abelicela, R. Plewa leg. et coll.; Greece, Crete, Lasithi pref., Lasithi plain, 24.-25.vi.1995, 1 spec., J. Batelka leg., P. Zahradnik coll.; Greece, Central Macedonia, Thessaloniki, Asprovalta, 15.v.2009, 1 spec., A. Márkus & T. Németh leg., HNHM coll.; Greece, Epirus, Preveza pref., Parga env., 28.vi.-6.vii.2000, 1 spec., S. Benedikt leg., P. Zahradnik coll.; Greece, Thessaly, Larisa pref., Olympos Mt., 15.v.2003, 1 spec., Z. Kraus leg. et coll.; Greece, South Aegean islands, Dodecanese pref., Rhodos I., Rhodos, vi.2009, 1 spec., A. Kotán leg., HNHM coll.; Greece, Thessaly, Trikala pref., Kalembara env., Kria Vrasi, Pindos Mt., 8.vi.2013, 1 spec., T. Sitek leg. et coll.

***Xestobium rufovillosum (DeGeer, 1774)***

*Xestobium rufovillosum* (DeGeer, 1774): Oertzen, 1886: 254; Yakobson, 1913: 817; Zahradnik, 2007a: 352.

*Anobium tessellatum* (Olivier, 1790): Kiesenwetter, 1859: 162.

**Distribution.** Europe, Palearctic Asia, introduced into North America and New Caledonia.

**Material examined.** Greece, Crete, Chania pref., Omalos, 21.v.2004, 2 spec., J. Januš leg. et coll.; Greece, Thessaly, Larisa pref., Stomio env., Ossa Oros Mt., 29.-31.v.2012, 2 spec., T. Sitek leg. et coll.

### **EUCRADINAE LECONTE, 1861** **Hedobiini Mulsant et Rey, 1868**

#### ***Hedobia pubescens* (Olivier, 1790)**

*Hedobia pubescens* (Olivier, 1790): Horion, 1961: 189; Burakowski, Mroczkowski & Stefańska, 1986: 47; Zahradník, 2007a: 353.

**Distribution.** Central and southern Europe, Caucasus, Turkey.

**Material examined.** Greece, Thessaly, Larisa pref., Thirio env., Akarnanika Mt., 28.v.1997, 1 spec. S. Benedikt leg., V. Benedikt coll.; Greece, Western Macedonia, Grevena pref., Karpero env., Anici, 5.-6.vi.2012, 2 spec., T. Sitek leg. et coll.

#### ***Ptinomorphus imperialis* (Linnaeus, 1767)**

*Hedobia imperialis* (Linnaeus, 1767): Horion, 1961: 191.

*Ptinomorphus imperialis* (Linnaeus, 1767): Español, 1977: 5; Zahradník, 2007a: 354.

**Distribution.** Europe, Caucasus, Turkey.

**Material examined.** Greece, Eastern Macedonia and Thrace, Drama pref., Livadero, Falakro Mt. Area, 10.v.2011, 1 spec., Z. Malinka leg. et coll.; Greece, Thessaly, Larisa pref., Spilia env., Ossa Oros Mt., 14.-16.vi.2010, 1 spec., J. Plecháč leg. et coll.; Greece, Thessaly, Larisa pref., Thirio env., Akarnanika Mt., 28.v.1997, 1 spec., S. Benedikt leg., V. Benedikt coll.

#### ***Ptinomorphus magnificus magnificus* (Reitter in Leder, 1880)**

*Ptinomorphus magnificus magnificus* (Reitter in Leder, 1880): Logvinovskiy, 1985: 40; Zahradník, 1998: 264; Toskina, 2001: 150; Zahradník, 2007a: 354.

*Hedobia magnifica* Reitter in Leder, 1880: Brenske et Reitter, 1884: 86; Oertzen, 1886: 254; Seidlitz, 1889: 497; Seidlitz, 1891: 531; Heyden, Reitter & Weise, 1891: 231; Heyden, Reitter & Weise, 1906: 427; Pic, 1912b: 6; Yakobson, 1913: 818; Schaufuss, 1916: 722; Zaytsev, 1956: 52.

**Distribution.** Caucasus, southern Balkan Peninsula, Turkey.

**Material examined.** Greece, Central Greece, Fthiotida pref., Aranova, Parnassus Mt., 5.-9.vi.1989, 1 spec., Kudrna leg., P. Zahradník coll.; Greece, Crete, Chania pref., Imbros, 20.v.2007, 1 spec., O. Konvička leg. et coll.; Greece, Crete, Chania pref., Samaria, 6.vi.1987, 1 spec., Kudrna leg., P. Zahradník coll.; Greece, Thessaly, Trikala pref., Neraida, 1.vi.2013, 1 spec., E. Ezer leg. et coll.

#### ***Ptinomorphus perpulchrus* (Obenberger, 1917)**

*Ptinomorphus perpulchrus* (Obenberger, 1917): Zahradník, 2007a: 354.

*Hedobia perpulchra* Obenberger, 1917: Obenberger, 1917: 34.

**Distribution.** Greece.

#### ***Ptinomorphus regalis* (Duftschmid, 1825)**

*Hedobia regalis* (Duftschmid, 1825): Horion, 1961: 190.

*Ptinomorphus regalis* (Duftschmid, 1825): Zahradník, 2007a: 354.

**Distribution.** Europa, Caucasus, Iran, Turkey.

***Ptinomorphus tatjanae* Logvinovskiy, 1978**

**Distribution.** Kyrgyzstan, Russia (South European Territory), Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Fourka env., Smolikas Mt., 5.vi.2008, 1 spec., K. Orszulik leg. et coll. **New species for Greece.**

**GIBBINAE JACQUELIN DU VAL, 1860****Gibbiini Jacquelin du Val, 1860*****Gibbium aequinoctiale* Boieldieu, 1854**

*Gibbium aequinoctiale* Boieldieu, 1854: Bellés, 1984: 7; Bellés, 1985: 41; Bellés & Mifsud, 2000: 46; Borowski, 2007: 329.

**Distribution.** Cosmopolitan due to introduction, original distribution unknown.

***Gibbium boieldieui* Levrat, 1857**

*Gibbium boieldieui* Levrat, 1857: Pic, 1912a: 5; Yakobson, 1913: 824; Bellés, 1985: 43; Borowski, 1996: 12; Borowski, 2007: 329.

**Distribution.** Caucasus and neighbouring countries, Greece.

***Gibbium psylloides* (Czempinski, 1778)**

*Gibbium psylloides* Czempinski, 1778: Oertzen, 1886: 253; Sahlberg, 1913: 17; Yakobson, 1913: 824; Bellés, 1984: 7; Bellés, 1985: 39; Buchelos, 1985: 224; Borowski, 2007: 329; Denux & Zagatti, 2010: 343.

*Gibbium psylloides* var. *brevicornis* Reitter, 1885: Heyden, Reitter & Weise, 1891: 228; Heyden, Reitter & Weise, 1906: 423; Pic, 1912a: 5.

*Gibbium scotias* (Fabricius, 1781): Kiesenwetter, 1859: 162.

**Distribution.** Europe, northern Africa, northern Asia. Introduced into China and certain European countries.

**Meziini Bellés, 1985*****Mezium affine* Boieldieu, 1856**

*Mezium affine* Boieldieu, 1856: Kiesenwetter, 1859: 162; Mulsant & Rey, 1868: 214; Oertzen, 1886: 253; Yakobson, 1913: 824; Horion, 1961: 258; Pelekassis, 1962: 67; Bellés, 1984: 13; Bellés, 1985: 50; Buchelos, 1985: 224; Borowski, 2007: 329.

**Distribution.** Central and southern Europe. Introduced into further European countries.

***Mezium americanum* (Castelnau, 1840)**

*Mezium americanum* (Castelnau, 1840): Reitter, 1884b: 297; Marseul, 1886: 301; Heyden, Reitter & Weise, 1891: 228; Heyden, Reitter & Weise, 1906: 423; Fauvel, 1904: 148; Fall, 1905: 103; Pic, 1912a: 6; Yakobson, 1913: 824; Schafuss, 1916: 716; Bellés, 1985: 53.

*Mezium arachnoides* Desbrochers des Loges, 1875: Desbrochers des Loges, 1875: 50; Stein & Weise, 1877: 107; Heyden, Reitter & Weise, 1883: 121; Oertzen, 1886: 253; Pic, 1912a: 6.

**Distribution.** Cosmopolitan, introduced into a number of European countries.

**MESOCOLEPODINAE MULSANT ET REY, 1864****Mesocoelopodini Mulsant et Rey, 1864**

**Mesocoelopus brevistriatus** Leiler, 1979

*Mesocoelopus brevistriatus* Leiler, 1979: Leiler, 1979: 95; Leiler, 1985: 143; Zahradník, 1998: 278; Zahradník, 2007a: 354.

**Distribution.** Greece, Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Geraxini, 7.x.1977, 1 spec., T.-E. Leiler leg., P. Zahradník coll. (paratype).

**Mesocoelopus creticus** Fairmaire, 1880

*Mesocoelopus creticus* Fairmaire, 1880: Fairmaire, 1880: 241; Marseul, 1886: 297; Oertzen, 1886: 255; Heyden, Reitter & Weise, 1891: 233; Schilsky, 1899: 37; Reitter, 1901: 39; Heyden, Reitter & Weise, 1906: 432; Pic, 1912b: 60; Yakobson, 1913: 810; Schaufuss, 1916: 727; Españaol, 1967: 101; Españaol, 1979a: 34; Zahradník, 2007a: 354.

**Distribution.** Greece.

**Material examined.** Greece, Crete, Chania pref., Omolos, 22.-30.v.1990, 2 spec., I. Jeniš leg., P. Zahradník coll.; Greece, South Aegean insels, Dodecanese pref., Rhodos I., Kolymbia, 23.-28.vi.2014, 8 spec., T. Staněk leg., T. Staněk et P. Zahradník coll.; Greece, Western Greece, Achaia pref., Diakopto, Vouraikos, vii.1999, 1 spec., ex larvae, B. Büche leg. et coll.

**Mesocoelopus niger** (P.W.J. Müller, 1821)

*Mesocoelopus niger* (P.W.J. Müller, 1821): Oertzen, 1886: 255; Schilsky, 1899: 39; Yakobson, 1913: 810; Horion, 1961: 243; Logvinovskiy, 1985: 141; Españaol, 1992: 136; Zahradník, 2007a: 354.

**Distribution.** Central and southern Europe, northern Africa, Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Crete, Rethymno pref., Plakias, 23.-25.vi.2005, 1 spec., Z. Švec leg. et coll.; Greece, Thessaly, Larisa pref., Olympiada, 6.vi.1999, 1 spec., Z. Švec leg. et coll.; Greece, Thessaly, Larisa pref., Olympiada, vi.2008, 10 spec., ex larvae from *Hedera helix*, Z. Kraus leg., Z. Kraus et P. Zahradník coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 24.v.2011, 1 spec., A. Kotán, A. Márkus & T. Németh leg., HNHM coll.; Greece, Thessaly, Larisa pref., Stomio env., Ossa Oros Mt., 1.vi.2011, 1 spec., O. Konvička leg. et coll.; Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 4.-6.vi.2013, 1 spec., T. Sitek leg. et coll.

**Mesoths ferrugineus** (Mulsant et Rey, 1861)

*Mesoths ferrugineus* (Mulsant et Rey, 1861): Schaufuss, 1916: 727; Horion, 1961: 243; Gottwald, 1975: 146; Zahradník, 2007a: 355.

**Distribution.** Southeastern Europe, Middle East, Turkey.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Vrasna, Asprovalta env., 8.-10.vi.2010, 1 spec., T. Sitek leg. et coll.; Greece, Eastern Macedonia and Thrace, Rodopi pref., Nea Santa, Mt. Sapka, 11.vi.2008, 2 spec., A. Kotán leg., HNHM coll.; Greece, Epirus, Preveza pref., Neochori env., Agia Apostoli, 8.vi.2008, 1 spec., B. Zbuzek leg., P. Zahradník coll.; Greece, Epirus, Preveza pref., Preveza env., vii.1991, 1 spec., O. Kapler leg., P. Zahradník coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kastraki, 3.-6.vi.2007, 2 spec., T. Sitek leg., T. Sitek et P. Zahradník coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kria Vrasi, 16.vi.2014, 1 spec., Z. Malinka leg. et coll.; Greece, Western Macedonia, Grevena pref., Karpero env., Anixi, 10.vi.2013, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kastraki, 3.-6.vi.2007, 2 spec., 11.-12.vi.2010, 6 spec., T. Sitek leg., T. Sitek et P. Zahradník coll.; Greece, Thessaly, Trikala pref., Kalambaka env., Kria Vrasi, 12.vi.2010, 4 spec., T. Sitek leg., T. Sitek et P. Zahradník coll.; Greece, Thessaly, Trikala pref., Trigona env., Kalambaka, 7.vi.2008, 1 spec., B. Zbuzek leg. et coll.; Greece, Western Greece, Achaia pref., Metochi, Patra env., 2.vi.2010, 5 spec., K. Rébl leg. et coll..

**PTILININAE SHUCKARD, 1839****Ptilinini** Shuckard, 1839

***Ptilinus fuscus* (Geoffroy in Fourcroy, 1785)**

*Ptilinus fuscus* (Geoffroy in Fourcroy, 1785): Zahradník, 2007a: 356.

**Distribution.** Europe, northern Africa, northern Asia, Japan.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Stavros, 1.-11.vi.1993, 1 spec., P. Vonička leg., R. Čtvrtečka coll.

**PTININAE LATREILLE, 1802****Ptinnini Latreille, 1802*****Dignomus atticus* (Pic, 1902)**

*Dignomus atticus* (Pic, 1902): Bellés, 1996: 221;

*Pinus* (*Eutaphrus*) *atticus* Pic, 1902: Pic, 1902b: 32; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 17; Yakobson, 1913: 822; Schaufuss, 1916: 719; Borowski, 2002: 36; Borowski, 2007: 330.

**Distribution.** Greece, Turkey.

***Dignomus aureopilis* (Desbrochers des Loges, 1875)**

*Dignomus aureopilis* (Desbrochers des Loges, 1875): Borowski, 2007: 330.

*Pinus* (*Eutaphrus*) *aureopilis* Desbrochers des Loges, 1875: Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 18; Yakobson, 1913: 822; Schaufuss, 1916: 719.

*Bruchus* (*Eutaphrus*) *aureopilis* (Desbrochers des Loges, 1875): Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Middle East, Egypt.

**Note:** According to Borowski (2007) this species does not occur in Greece.

***Dignomus canaliculatus* (Reitter in Brenske et Reitter, 1884)**

*Dignomus canaliculatus* (Reitter in Brenske et Reitter, 1884): Bellés, 1996: 220; Borowski, 2002: 36; Borowski, 2007: 330.

*Pinus* (*Eutaphrus*) *canaliculatus* Reitter in Brenske et Reitter, 1884: Marseul, 1886: 299; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 19; Yakobson, 1913: 822; Schaufuss, 1916: 719.

*Dignomus graecus* (Pic, 1896): Bellés, 1996: 221.

*Pinus* (*Eutaphrus*) *graecus* (Pic, 1896): Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 24; Yakobson, 1913: 822; Schaufuss, 1916: 719.

*Eutaphrus graecus* Pic, 1896: Pic, 1896: 41.

*Bruchus* (*Eutaphrus*) *canaliculatus* Reitter in Brenske et Reitter, 1884: Brenske et Reitter, 1884: 85; Reitter, 1884b: 319; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 230.

**Distribution.** Greece.

**Material examined.** Greece, Crete, Lasithi pref., Vai Palleetrend, 1 spec., Pülscher leg., P. Zahradník coll.

***Dignomus frivaldszkyi* (Reitter, 1884)**

*Dignomus frivaldszkyi* (Reitter, 1884): Bellés, 1996: 217; Borowski, 2002: 37; Borowski, 2007: 330.

*Pinus* (*Eutaphrus*) *frivaldszkyi* (Reitter, 1884): Marseul, 1886: 299; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1906: 426; Sahlberg, 1913: 17; Pic, 1912a: 22; Yakobson, 1913: 822; Schaufuss, 1916: 719.

*Bruchus* (*Eutaphrus*) *frivaldszkyi* Reitter, 1884: Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Middle East, Cyprus, Greece, Turkey.

**Material examined.** Greece, Central Greece, Fthiotida pref., Aranova, Parnassus Mt., 5.v.1994, 1 spec., I. Jeniš leg., P. Zahradník coll.; Greece, Crete, Chania pref., Laki, 7.-19.v.2003, 8 spec., P. Moravec leg., P. Zahradník coll.; Greece, Crete, Chania pref., Omalos, 1.vi.2007, 1 spec., Z. Švec leg. et coll.; Greece, Crete, Chania pref., Sfakia, 7.-19.v.2003, 1 spec., P. Moravec leg., P. Zahradník coll.; Greece, Crete, Lasithi pref., Pantanassos, Reissler leg., J. Háva coll.; Greece, Epirus, Preveza pref., Preveza, 6.-16.vi.1992, 6 spec., T. Růžicka leg., T. Růžicka et P. Zahradník coll.; Greece, Epirus,

Thesprotia pref., Sivota, 20.-27.v.2000, 1 spec., F. Kantner leg., USNM coll.; Greece, Peloponnese, Arcadia pref., Tripoli env., 4.vi.2009, 1 spec., T. Růžička leg. et coll.; Greece, Peloponnese, Lakonia pref., Artemisia, Taygetos Mt., 1.-2.vi.2006, 1 spec., F. Houška leg. et coll.; Greece, Thessaly, Magnisia pref., Volos, Pelio Mt., 3.vi.2011, 1 spec., T. Staněk leg. et coll.; Greece, Western Macedonia, Grevena pref., Karpero env., Anixi, 18.vi.2014, 1 spec., Z. Malinka leg. et coll.

### ***Dignomus irroratus* (Kiesenwetter, 1851)**

*Dignomus irroratus* (Kiesenwetter, 1851): Borowski, 1993: 203; Borowski, 1996: 24; Borowski, 2002: 37; Borowski, 2007: 330.

*Pinus* (*Eutaphrus*) *irroratus* Kiesenwetter, 1851: Kiesenwetter, 1867: 129; Jakobson, 1913: 822.

*Bruchus irroratus* (Kiesenwetter, 1851): Oertzen, 1886: 254.

*Pinus alpinus* Boieldieu, 1854: Boieldieu, 1854: lxxviii; Kiesenwetter, 1859: 162.

**Distribution.** Southern Europe, introduced into Egypt, Germany, Poland.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Crete, Lasithi pref., Ellounda, 16.x.1969, 1 spec., Reissler leg., P. Zahradník coll.; Greece, South Aegean islands, Cyclades pref., Paros I., Naoussa, vii.1997, 1 spec., ex larvae from goat dung, B. Büche leg. et coll.

### ***Dignomus jacqueti* (Pic, 1895)**

**Distribution.** France, Italy, Spain.

**Material examined.** Greece, Calab, 2 spec., Paganetti leg., P. Zahradník coll. **New species for Greece.**

**Note:** Exact location is impossible. May be that there is a mistake. The specimens are labelled „Greece, Calab“, but „Calab“ may be abbreviation for Calabria in Italy. Possibly a species uncertain for Greece.

### ***Dignomus lesinae* (Reitter in Brenske et Reitter, 1884)**

*Pinus* (*Eutaphrus*) *lesinae* Reitter in Brenske et Reitter, 1884: Jakobson, 1913: 822.

*Bruchus lesinae* (Reitter in Brenske et Reitter, 1884): Oertzen, 1886: 254.

**Distribution.** Croatia.

**Note:** According to Borowski (2007) this species does not occur in Greece.

### ***Dignomus quadricornis* (Pic, 1895)**

*Dignomus quadricornis* (Pic, 1895): Bellés, 1996: 211; Borowski, 2002: 37; Borowski, 2007: 331.

*Pinus* (*Eutaphrus*) *quadricornis* (Pic, 1895): Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 31; Jakobson, 1913: 823; Schaufuss, 1916: 719;

**Distribution.** Middle East, Cyprus, Greece, Turkey.

### ***Dignomus reichei* (Boieldieu, 1854)**

*Dignomus reichei* (Boieldieu, 1854): Bellés, 1996: 224; Bellés & Mifsud, 2000: 43; Borowski, 2002: 37; Borowski, 2007: 331.

*Pinus* (*Eutaphrus*) *reichei* Boieldieu, 1854: Boieldieu, 1854: lxxix; Kiesenwetter, 1859: 162; Mulsant & Rey, 1868: 65; Kiesenwetter, 1877: 52; Stein & Weise, 1877: 107; Marseul, 1886: 299; Sahlberg, 1903b: 63; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 32; Jakobson, 1913: 823; Schaufuss, 1916: 719; Novak, 1964: 88.

*Bruchus* (*Eutaphrus*) *reichei* (Boieldieu, 1854): Reitter, 1884b: 320; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 230.

*Dignomus monstrositarsis* (Pic, 1896): Bellés, 1996: 225.

**Distribution.** Albania, Croatia, Greece.

**Material examined.** Greece, Epirus, Arta pref., Petra env., 7.vi.2005, 1 spec., J. Plecháč leg., P. Zahradník coll.; Greece, Ionian Islands, Korfu pref., Ag. Georgios, L. Korission, 19.-27.vi.2003, 1 spec., K. Schön leg., P. Zahradník coll.; Greece, Ionian Islands, Korfu pref., Korfu I., 1 spec., U. Sahlberg leg., P. Zahradník coll.; Greece, Peloponnese, Lakonia pref., Artemisia, Taygetos Mt., 1.-2.vi.2006, 1 spec., F. Houška leg. et coll.

***Dignomus xylopertha* (Boieldieu, 1856)**

*Dignomus xylopertha* (Boieldieu, 1856): Borowski, 2007: 331.

*Ptinus xylopertha* Boieldieu, 1856: Heyden, Reitter & Weise, 1906: 426; Schaufuss, 1916: 719.

*Bruchus* (*Eutaphrus*) *xylopertha* (Boieldieu, 1856): Reitter, 1884b: 320; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 230.

*Ptinus* (*Eutaphrus*) *xylopertha* Reiche et Saulcy, 1857: Pic, 1912a: 37; Yakobson, 1913: 822.

**Distribution.** Northern Africa, Middle East, Cyprus, Greece, Turkey.

***Epauloecus unicolor* (Piller et Mitterpacher, 1783)**

*Epauloecus unicolor* (Piller et Mitterpacher, 1783): Borowski, 2007: 331.

**Distribution.** Europe, Caucasus, introduced into North America.

***Eurostodes quisquilarium* (Baudi di Selve, 1874)**

*Eurostodes quisquilarium* (Baudi di Selve, 1874): Schaufuss, 1916: 717.

**Distribution.** Italy.

**Note:** According to Borowski (2007) this species does not occur in Greece.

***Niptus* (*Niptus*) *hololeucus* (Faldermann, 1835)**

*Niptus* (*Niptus*) *hololeucus* (Faldermann, 1835): Borowski, 2007: 332.

**Distribution.** Cosmopolitan.

***Ptinus* (*Bruchoptinus*) *brevivittis* Reitter, 1881**

*Ptinus* (*Bruchoptinus*) *brevivittis* Reitter, 1881: Yakobson, 1913: 820; Borowski, 2007: 333.

*Bruchus* (*Gynopteris*) *brevivittis* (Reitter, 1881): Brenske et Reiter, 1884: 83; Reitter, 1884b: 307; Oertzen, 1886: 254.

**Distribution.** Balkan Peninsula.

***Ptinus* (*Bruchoptinus*) *femoralis* (Reitter, 1884)**

*Ptinus* (*Bruchoptinus*) *femoralis femoralis* (Reitter, 1884): Heyden, Reitter & Weise, 1906: 425; Yakobson, 1913: 821; Schaufuss, 1916: 719; Iablokoff-Khznorian & Karapetyan, 1991: 24; Borowski, 2007: 333.

*Bruchus* (*Bruchoptinus*) *femoralis* Reitter, 1884: Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Algeria, Cyprus, Greece, Italy, Spain, Tunisia.

**Material examined.** Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 3 spec., Z. Švec leg.; Greece, Thessaly, Trikala pref., Kalambaka env., Kria Vrisi, 16.vi.2014, 1 spec., Z. Malinka leg. et coll.

***Ptinus* (*Bruchoptinus*) *henoni* Pic, 1897**

*Ptinus* (*Bruchoptinus*) *henoni* Pic, 1897: Borowski, 2007: 333.

**Distribution.** Azerbaijan, Greece, Turkey.

**Material examined.** Greece, Thessaly, Larisa pref., Thirio env., Akarnanika Mt., 28.v.1997, 2 spec., S. Benedikt leg., V. Benedikt coll.

***Ptinus (Bruchoptinus) pellitus* Desbrochers des Loges, 1875**

*Ptinus (Bruchoptinus) pellitus* Desbrochers des Loges, 1875: Borowski, 2007: 333.

**Distribution.** Cyprus, Greece (Crete), Lebanon, Syria, Turkey.

**Material examined.** Greece, Crete, Chania pref., Elos, 6.-20.v.2003, 1 spec., P. Viktora leg., P. Zahradník coll.; Greece, Crete, Chania pref., Omalos, 4.vi.2011, 1 spec., Z. Švec leg. et coll.

***Ptinus (Bruchoptinus) rufipes* Olivier, 1790**

*Ptinus (Bruchoptinus) rufipes* Olivier, 1790: Borowski, 2007: 333.

*Ptinus (Bruchoptinus) brunnescens* Pic, 1896: Pic, 1912a: 19; Yakobson, 1913: 820.

**Distribution.** Europe.

**Material examined.** Greece, Central Macedonia, Pieria pref., Litochoro, 6.vi.2003, 1 spec., R. Škoda leg. et coll.; Greece, Central Macedonia, Pieria pref., Priona-Litochoro, Olympos Mt., 3.vi.2003, 1 spec., R. Škoda leg. et coll.

***Ptinus (Bruchoptinus) schatzmayeri* Pic, 1934**

*Ptinus (Bruchoptinus) schatzmayeri* Pic, 1934: Pic, 1934: 29.

**Distribution.** Greece.

**Note:** Pic (1934) described this species from Mont Athos, Macedonia. But he intended the province in Greece of this name, not the country Macedonia (formerly a part of Yugoslavia). Borowski (2007) quoted this species by mistake not from Greece, but from Macedonia.

***Ptinus (Bruchoptinus) torretassoii* Pic, 1934**

*Ptinus (Bruchoptinus) torretassoii* Pic, 1934: Pic, 1934: 29; Borowski, 2007: 333.

**Distribution.** Greece, Turkey.

**Material examined.** Greece, Epirus, Thesprotia pref., Sivota, 85 km S of Igoumenitsa, 20.-27.v.2000, 1 spec., F. Kantner leg., P. Zahradník coll.

***Ptinus (Cyphoderes) bidens* Olivier, 1790**

*Ptinus (Cyphoderes) bidens* Olivier, 1790: Horion, 1961: 282; Bellés, 2002: 99; Borowski, 2007: 333.

*Bruchus bidens* (Olivier, 1790: Oertzen, 1886: 254.

**Distribution.** Southern and western Europe, introduced to Sweden and the Canary Islands.

***Ptinus (Cyphoderes) raptor* (Sturm, 1837)**

*Ptinus (Cyphoderes) raptor* (Sturm, 1837): Borowski, 2007: 333.

**Distribution.** Europe, Siberia, Kazakhstan, Mongolia, introduced into North America.

***Ptinus (Cyphoderes) schlerethi* (Reitter, 1884)**

*Ptinus (Cyphoderes) schlerethi* (Reitter, 1884): Marseul, 1886: 300; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 33; Yakobson, 1913: 822; Schaufuss, 1916: 719; Horion, 1961: 281; Iablokoff-Khnzorian & Karapetyan, 1991: 53; Bellés, 2002: 102; Borowski, 2007: 333.

*Bruchus (Cyphoderes) schlerethi* Reitter, 1884: Reitter, 1884b: 317; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Central and southeastern Europe.

**Material examined.** Greece, Central Macedonia, Pieria pref., Karitsa, Kissavoss Mt., 13.iv.2006, 1 spec., S. Vít leg., P. Zahradník coll.



***Ptinus (Gynopterus) crassicornis* Kiesenwetter, 1877**

*Ptinus (Gynopterus) crassicornis* Kiesenwetter, 1877: Marseul, 1886: 299; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 20; Jakobson, 1913: 822; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 19; Borowski, 2007: 334.

*Bruchus crassicornis* (Kiesenwetter, 1877): Reitter, 1884b: 315; Oertzen, 1886: 253.

*Bruchus forticornis* (Reitter, 1878): Brenske et Reiter, 1884: 83.

**Distribution.** Greece, Italy.

***Ptinus (Gynopterus) dubius* Sturm, 1837**

*Ptinus (Gynopterus) dubius* Sturm, 1837: Jakobson, 1913: 822; Horion, 1961: 275; Borowski, 2007: 334.

*Bruchus dubius* (Sturm, 1837): Oertzen, 1886: 254.

**Distribution.** Europe, northern Africa, Turkey.

***Ptinus (Gynopterus) salvatori* Pic, 1901**

*Ptinus (Gynopterus) salvatori* Pic, 1901: Pic, 1901: 52; Heyden, Reitter & Weise, 1906: 425; Pic, 1912a: 33; Jakobson, 1913: 821; Schaufuss, 1916: 719; Borowski, 2007: 334.

**Distribution.** Greece.

***Ptinus (Gynopterus) sexpunctatus* Panzer, 1789**

*Ptinus (Gynopterus) sexpunctatus* Panzer, 1789: Borowski, 2007: 334.

**Distribution.** Europe.

***Ptinus (Gynopterus) subroseus* Reitter, 1888**

*Ptinus (Gynopterus) subroseus* Reitter, 1888: Fauvel, 1889: 183; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 35; Jakobson, 1913: 822; Schaufuss, 1916: 719; Borowski, 2007: 334.

*Bruchus (Gynopterus) subroseus* (Reitter, 1888): Oertzen, 1886: 254 (nom. nud.); Reitter, 1888: 430; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Greece.

***Ptinus (Gynopterus) variegatus* Rossi, 1794**

*Ptinus (Gynopterus) variegatus* Rossi, 1794: Kiesenwetter, 1859: 162; Sahlberg, 1903b: 63; Jakobson, 1913: 822; Horion, 1961: 278; Borowski, 2007: 334.

*Bruchus variegatus* (Rossi, 1794): Oertzen, 1886: 253.

**Distribution.** Europe, northern Africa, Middle East, introduced into North America.

***Ptinus (Pseudoptinus) cumaniensis* Pic, 1896**

*Ptinus (Pseudoptinus) cumaniensis* Pic, 1896: Heyden, Reitter & Weise, 1906: 424; Pic, 1912a: 20; Jakobson, 1913: 820; Schaufuss, 1916: 719; Borowski, 2007: 334.

**Distribution.** Greece.

**Material examined.** Greece, Western Greece, Achaia, Kalavryta, Aorania, 30.iii.1997, 1 spec., L. Zerche leg., P. Zahradnik coll.

***Ptinus (Pseudoptinus) nikitanus (Brenske et Reitter, 1884)***

**Distribution.** Croatia, Yugoslavia (Serbia or Monte Negro, or both).

**Material examined.** Greece, Central Macedonia, Pieria pref., Pieria, Ritini, Mt., 18.iv.2006, 1 spec., S. Vít leg., P. Zahradník coll. **New species for Greece.**

***Ptinus (Pseudoptinus) oertzeni Reitter, 1888***

*Ptinus (Pseudoptinus) oertzeni* Reitter, 1888: Fauvel, 1889: 183; Heyden, Reitter & Weise, 1906: 424; Pic, 1912a: 29; Jakobson, 1913: 820; Schaufuss, 1916: 719; Borowski, 2007: 335.

*Ptinus (Pseudoptinus) leucaspis* Reitter, 1888: Fauvel, 1889: 183; Heyden, Reitter & Weise, 1906: 424; Pic, 1912a: 27; Jakobson, 1913: 820; Schaufuss, 1916: 719.

*Bruchus (Pseudoptinus) oertzeni* (Reitter, 1888): Oertzen, 1886: 254 (nom. nud.); Reitter, 1888: 428; Heyden, Reitter & Weise, 1891: 229.

*Bruchus (Pseudoptinus) leucaspis* (Reitter, 1888): Reitter, 1888: 428; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Greece.

***Ptinus (Pseudoptinus) rugosicollis Desbrochers des Loges, 1875***

*Ptinus (Pseudoptinus) rugosicollis* Desbrochers des Loges, 1875: Borowski, 2007: 335.

**Distribution.** Northern Africa (western part), Middle East, Cyprus, Greece, Turkey.

***Ptinus (Pseudoptinus) subaeneus (Reitter in Brenske et Reitter, 1884)***

*Ptinus (Pseudoptinus) subaeneus* (Reitter in Brenske et Reitter, 1884): Heyden, Reitter & Weise, 1906: 424; Sahlberg, 1913: 17; Pic, 1912a: 35; Jakobson, 1913: 820; Schaufuss, 1916: 719; Iablokoff-Khznorian & Karapetyan, 1991: 28; Borowski, 2007: 335.

*Bruchus (Pseudoptinus) subaeneus* Reitter in Brenske et Reitter, 1884: Brenske et Reitter, 1884: 81; Reitter, 1884b: 304; Oertzen, 1886: 253; Heyden, Reitter & Weise, 1891: 228.

*Ptinus (Pseudoptinus) pistaciae* Pic, 1901: Pic, 1901: 51; Heyden, Reitter & Weise, 1906: 424; Pic, 1912a: 31; Jakobson, 1913: 820; Schaufuss, 1916: 719.

**Distribution.** Middle East, Cyprus, Greece, Turkey.

**Material examined.** Greece, Peloponnese, Lakonia pref., Mavrovouni, Gythion, 17.iv.1979, 1 spec., S. Vít leg., P. Zahradník coll.

***Ptinus (Ptinus) argolisanus (Reitter, 1884)***

*Ptinus (Ptinus) argolisanus* (Reitter, 1884): Marseul, 1886: 300; Seidlitz, 1889: 509; Seidlitz, 1891: 545; Heyden, Reitter & Weise, 1906: 425; Pic, 1912a: 17; Jakobson, 1913: 821; Schaufuss, 1916: 719;

*Bruchus (Bruchus) argolisanus* Reitter, 1884: Reitter, 1884b: 310; Heyden, Reitter & Weise, 1891: 229; Borowski, 2007: 335.

**Distribution.** Greece.

***Ptinus (Ptinus) calcaratus Kiesenwetter, 1877***

*Ptinus (Ptinus) calcaratus* Kiesenwetter, 1877: Iablokoff-Khznorian & Karapetyan, 1991: 46; Borowski, 2007: 335.

*Ptinus (Ptinus) edmundi* Abeille de Perrin, 1894: Horion, 1961: 272.

*Ptinus (Ptinus) perplexus* Reitter, 1884: Sahlberg, 1903b: 63; Jakobson, 1913: 821.

*Bruchus perplexus* (Reitter, 1884): Oertzen, 1886: 254.

**Distribution.** Central and southern Europe.

**Material examined.** Greece, Eastern Macedonia and Thrace, Xanthi pref., Stavropouli, 25.-26.iv.2007, 2 spec., T. Růžička leg., T. Růžička et P. Zahradník coll.

***Ptinus (Ptinus) calcarifer* Reitter, 1888**

*Ptinus (Ptinus) calcarifer* Reitter, 1888: Fauvel, 1889: 183; Heyden, Reitter & Weise, 1906: 425; Pic, 1912a: 19; Jakobson, 1913: 822; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 50; Borowski, 2007: 335.  
*Bruchus (Heteroptinus) calcarifer* (Reitter, 1888): Reitter, 1888: 429; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Greece.

**Material examined.** Greece, Peleoponnese, Lakonia pref., Sparti Kalamata, Tripi, 8.iii.1995, 1 spec., Sabella leg., P. Zahradnik coll.

***Ptinus (Ptinus) fur* (Linnaeus, 1758)**

*Ptinus (Ptinus) fur* (Linnaeus, 1758): Jakobson, 1913: 821; Borowski, 2007: 336; Denux & Zagatti, 2010: 345.  
*Bruchus fur* (Linnaeus, 1758): Oertzen, 1886: 254.

**Distribution.** Cosmopolitan, originally Palearctic.

***Ptinus (Ptinus) gylippus* Reitter, 1906**

*Ptinus (Ptinus) gylippus* Reitter, 1906: Reitter, 1906: 283; Pic, 1912a: 24; Jakobson, 1913: 821; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 52; Borowski, 2007: 336.

**Distribution.** Greece.

***Ptinus (Ptinus) kiesenwetteri* (Reitter in Brenske et Reitter, 1884)**

*Ptinus (Ptinus) kiesenwetteri* (Reitter in Brenske et Reitter, 1884): Marseul, 1886: 300; Oertzen, 1886: 254; Sahlberg, 1903a: 8; Heyden, Reitter & Weise, 1906: 425; Pic, 1912a: 25; Jakobson, 1913: 822; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 50; Borowski, 2007: 336.

*Ptinus (Ptinus) kiesenwetteri* var. *cytherias* Pic, 1903: Pic, 1903: 170; Heyden, Reitter & Weise, 1906: 425; Schaufuss, 1916: 719.

*Bruchus (Heteroplus) kiesenwetteri* Reitter in Brenske et Reitter, 1884: Brenske et Reitter, 1884: 83; Reitter, 1884b: 314; Heyden, Reitter & Weise, 1891: 229.

*Ptinus (Ptinus) creticus* lablokoff-Khnzorian & Karapetyan, 1991: lablokoff-Khnzorian & Karapetyan, 1991: 50.

**Distribution.** Greece, Syria, Turkey.

***Ptinus (Ptinus) kruperi* Pic, 1929**

*Ptinus (Ptinus) kruperi* Pic, 1929: Pic, 1929: 14; Borowski, 2007: 336.

**Distribution.** Greece.

**Material examined.** Greece, Sanda, Mt. Sapka, Thrakia, 5.v.2009, 1 spec., P. Zahradnik coll.

***Ptinus (Ptinus) latro* Fabricius, 1775**

*Ptinus (Ptinus) latro* Fabricius, 1775: Jakobson, 1913: 821; Borowski, 2007: 336; Denux & Zagatti, 2010: 345.

*Ptinus (Ptinus) clavipes* Panzer, 1792: Buchelos, 1994: 102.

*Bruchus latro* (Fabricius, 1775): Oertzen, 1886: 254.

*Bruchus corticinus* Rottenberg, 1870: Brenske et Reiter, 1884: 83.

*Ptinus brunneus* Duftschmid, 1825: Kiesenwetter, 1859: 162; Sahlberg, 1903a: 3; Sahlberg, 1903b: 63.

*Bruchus brunneus* (Duftschmid, 1825): Oertzen, 1886: 254.

**Distribution.** Cosmopolitan, originally Palearctic.

***Ptinus (Ptinus) obesus* Lucas, 1849**

*Ptinus (Ptinus) obesus* Lucas, 1849: Jakobson, 1913: 821(?); lablokoff-Khnzorian & Karapetyan, 1991: 50; Borowski, 2007: 336.

*Bruchus obesus* (Lucas, 1849): Oertzen, 1886: 254.

**Distribution.** Sporadic in southern Europe, northern Africa.

***Ptinus (Ptinus) phlomidis* Boieldieu, 1854**

*Ptinus (Ptinus) phlomidis* Boieldieu, 1854: Boieldieu, 1854: lxxx; Kiesenwetter, 1859: 162; Gemminger & Harold, 1869: 1765; Stein & Weise, 1877: 107; Calwer, 1884: 376; Marseul, 1886: 300; Sahlberg, 1903b: 63; Heyden, Reitter & Weise, 1906: 425; Jakobson, 1913: 821; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 44; Borowski, 2007: 336.

*Bruchus (Bruchus) phlomidis* (Boieldieu, 1854): Reitter, 1884b: 314; Oertzen, 1886: 253; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Balkan Peninsula, Hungary, Italy, Turkey.

**Material examined.** Greece, Thessaly, Magnisia pref., Volos, Pelio Mt., 3.vi.2011, 1 spec., T. Staněk leg., P. Zahradník coll.

***Ptinus (Ptinus) pilosus* P.W.J. Müller, 1821**

*Ptinus (Ptinus) pilosus* P.W.J. Müller, 1821: Sahlberg, 1903b: 63; Jakobson, 1913: 821; Horion, 1961: 272; lablokoff-Khnzorian & Karapetyan, 1991: 49; Borowski, 2007: 337.

*Bruchus pilosus* (P.W.J. Müller, 1821): Oertzen, 1886: 254.

**Distribution.** Europe (except Pyrenean Peninsula and Russia except southern part).

***Ptinus (Ptinus) rufus* Brullé, 1832**

*Ptinus (Ptinus) rufus* Brullé, 1832: Kiesenwetter, 1859: 162; Gemminger & Harold, 1869: 1766; Stein & Weise, 1877: 107; Calwer, 1884: 376; Heyden, Reitter & Weise, 1906: 426; Pic, 1912a: 33; Jakobson, 1913: 823; Schaufuss, 1916: 719; Borowski, 2007: 337.

*Bruchus (Eutaphrus) rufus* (Brullé, 1832): Heyden, Reitter & Weise, 1891: 230.

**Distribution.** Greece.

***Ptinus (Ptinus) spitzyi* A. Villa et G.B. Villa, 1838**

*Ptinus (Ptinus) spitzyi* A. Villa et G.B. Villa, 1838: Horion, 1961: 273.

**Distribution.** Western Europe, northern Africa, Bosna Hercegovina, Croatia.

**Note:** According to Borowski (2007) this species does not occur in Greece.

***Ptinus (Ptinus) subpilosus* Sturm, 1837**

*Ptinus (Ptinus) subpilosus* Sturm, 1837: Jakobson, 1913: 821; Saalas, 1923: 186; Borowski, 2007: 337.

*Bruchus subpilosus* (Sturm, 1837): Oertzen, 1886: 254.

**Distribution.** Europe.

***Ptinus (Ptinus) tarsalis* (Reitter in Brenske et Reitter, 1884)**

*Ptinus (Ptinus) tarsalis* (Reitter in Brenske et Reitter, 1884): Marseul, 1886: 300; Heyden, Reitter & Weise, 1906: 425; Pic, 1912a: 35; Jakobson, 1913: 822; Schaufuss, 1916: 719; lablokoff-Khnzorian & Karapetyan, 1991: 48; Borowski, 2007: 337.

*Bruchus (Heteroplus) tarsalis* Reitter in Brenske et Reitter, 1884: Brenske et Reitter, 1884: 84; Reitter, 1884b: 314; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 229.

**Distribution.** Cyprus, Greece, Italy.

**Material examined.** Greece, South Aegean islands, Dodecanese pref., Karpathos I., Lastos Nordhang, Oros Kolla, 16.iv.1999, 1 spec., H. Meybohm leg., P. Zahradník coll.

### ***Ptinus (Ptinus) villiger* (Reitter, 1884)**

*Ptinus (Ptinus) villiger* (Reitter, 1884): Borowski, 2007: 337.

**Distribution.** Europe, Caucasus, Siberia, Far East, Central Asia, introduced into North America.

### ***Ptinus (Tectoptynus) exulans* Erichson, 1842**

*Ptinus (Tectoptynus) exulans* Erichson, 1842: Borowski, 2007: 337.

**Distribution.** Cosmopolitan, originally from Australia.

### ***Ptinus (Tectoptynus) tectus* Boieldieu, 1856**

*Ptinus (Tectoptynus) tectus* Boieldieu, 1856: Jakobson, 1913: 821; Borowski, 2007: 337; Denux & Zagatti, 2010: 346.

**Distribution.** Cosmopolitan, originally from Australia.

### **Sphaericini Portevin, 1931**

#### ***Sphaericus (Sphaericus) pinguis* (Boieldieu, 1854)**

*Sphaericus (Sphaericus) pinguis* (Boieldieu, 1854): Borowski, 2007: 338.

**Distribution.** Cosmopolitan, originally from western Mediterranean, Azores Islands, Canary Islands, Madeira Islands.

#### ***Trigonogenius globosus* Solier, 1849**

*Trigonogenius globosus* Solier, 1849: Borowski, 2007: 339.

**Distribution.** Cosmopolitan, originally from South America.

### **XYLETININAE GISTL, 1848 Lasiodermini Böving, 1927**

#### ***Lasioderma aterrimum* Roubal, 1916**

*Lasioderma aterrimum* Roubal, 1916: Zahradník, 2007a: 357.

**Distribution.** Croatia, Czech Republic, Germany, Greece, Slovakia, Russia (South European Territory).

**Material examined.** Greece: Epirus, Arta pref., Koronisia, 22.v.1997, 1 spec., S. Benedikt leg., V. Benedikt coll.

#### ***Lasioderma baudii* Schilsky, 1899**

*Lasioderma baudii* Schilsky, 1899: Zahradník, 2007a: 357.

**Distribution.** The Mediterranean area.

**Material examined.** Greece, Crete, Chania pref., Paralia env., Katherini, 1.-10.viii.1997, 3 spec., M. Mantič leg. et coll.; Greece, Epirus, Preveza pref., Preveza, 15.vi.1999, 30 spec., Z. Švec leg., Z. Švec et P. Zahradník coll.

### *Lasioderma bubalus phelipaearun* Peyerimhoff, 1926

**Distribution.** Algeria, Morocco, Tunisia.

**Material examined.** Greece, Peloponnese, Lakonia pref., Anoqia env., Taygetos Mt., 20.v.2007, 1 spec., K. Rébl leg., P. Zahradník coll. **New species for Greece.**

### *Lasioderma cretaeum* Toskina, 2013

*Lasioderma cretaeum* Toskina, 2013: Toskina, 2013: 6.

**Distribution.** Greece.

### *Lasioderma fuscum* (Rey, 1892)

*Lasioderma fuscum* (Rey, 1892): Pic, 1912b: 55; Yakobson, 1913: 812; Schaufuss, 1916: 726; Zahradník, 2007a: 357. *Pseudochina fusca* Rey, 1892: Rey, 1892: 3.

**Distribution.** France, Greece.

### *Lasioderma haemorrhoidale* (Illiger, 1807)

*Lasioderma haemorrhoidale* (Illiger, 1807): Oertzen, 1886: 254; Schilsky, 1899: 33; Reitter, 1901: 34; Heyden, Reitter & Weise, 1906: 431; Pic, 1912b: 55; Yakobson, 1913: 812; Buchelos, 1989: 9; White, 1990: 346; Zahradník, 2007a: 257.

*Lasioderma haemorrhoidale* var. *unicolor* Schilsky, 1899: Schaufuss, 1916: 726.

**Distribution.** The Mediterranean area.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Rentina, 11.-30.vii.2004, 1 spec., Z. Kraus leg. et coll.; Greece, Crete, Chania pref., Paralia env., Katherini, 1.-10.viii.1998, 2 spec., M. Mantič leg. et coll.; Greece, Crete, Rethymno pref., Plakias env., 6.-16.vii.2005, 1 spec., P. Zahradník leg. et coll.

### *Lasioderma kiesenwetteri* Schilsky, 1899

*Lasioderma kiesenwetteri* Schilsky, 1899: Schilsky, 1899: 31; Reitter, 1901: 36; Heyden, Reitter & Weise, 1906: 431; Pic, 1912b: 56; Yakobson, 1913: 812; Schaufuss, 1916: 727; Horion, 1961: 243; Logvinovskiy, 1977b: 286; Logvinovskiy, 1985: 135; Zahradník, 2007a: 357.

**Distribution.** Southern Europe, occasionally appears in central Europe, Caucasus, Middle East, Central Asia.

**Material examined.** Greece, Crete, Chania pref., Agia Marina, 15.-24.vi.2004, 1 spec., I. Daněk leg. et coll.; Greece, Crete, 17.v.1975, 1 spec., USNM coll.; Greece, Eastern Macedonia and Thrace, Xanthi pref., Pezoula env., Neraida, Pindos, 9.vi.2013, 2 spec., T. Sitek leg. et coll.; Greece, Ionian Islands, Korfu pref., Potamos, v.1964, 1 spec., T. Palm leg., P. Zahradník coll.; Greece, South Aegean islands, Cyclades pref., Paros I., Naoussa, vii.19997, 2 spec., ex larvae from stumps of *Ferula communis*, B. Büche leg. et coll.

### *Lasioderma laeve* (Illiger, 1807)

*Lasioderma laeve* (Illiger, 1807): Oertzen, 1886: 255.

**Distribution.** Austria, France, Hungary, Italia, Portugal, Spain, Switzerland.

**Material examined.** Greece, Crete, Lasithi pref., Mochos, 11.vii.2013, 2 spec., T. Gazurek leg. et coll.

***Lasioderma nigellum* Toskina, 2013**

*Lasioderma nigellum* Toskina, 2013: Toskina, 2013: 15.

**Distribution.** Greece.

***Lasioderma obscurum* (Solsky, 1867)**

*Lasioderma obscurum* (Solsky, 1867): Pic, 1912b: 56; Horion, 1961: 243; Gottwald, 1970: 123; Emets, 1976: 194; Logvinovskiy, 1985: 137; Zahradník, 2007a: 358.

**Distribution.** Southern Europe, occasionally appears in central Europe, Central Europe, southeastern Russia.

**Material examined.** Greece, Central Greece, Fthiotida pref., Karpenissi env., Timfristos Mt., 26.vi.2005, 1 spec., S. Benedikt leg., P. Zahradník coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.-20.vii.2002, 1 spec., Z. Kraus leg. et coll.; Greece, Crete, Chania pref., Agia Marina env., 15.-24.vi.2003, 2 spec., L. Daněk leg. et coll.; Greece, Crete, Chania pref., Agia Marina, Litanakis, 1.vi.2004, 1 spec., J. Januš leg. et coll.; Greece, Crete, Chania pref., Kournas, 23.v.2004, 1 spec., J. Januš leg. et coll.; Greece, Crete, Chania pref., Omalos, 17.v.1994, 1 spec., S. Lundbreg leg. et coll.; Greece, Epirus, Thesprotia pref., Igoumenitsa, 1.vi.2001, 2 spec., R. Röber leg. et coll.; Greece, Ionian Islands, Zakhyntos pref., Argasi env., 5.-14.v.2013, 1 spec., T. Staněk leg. et coll.; Greece, Thessaly, Larisa pref., Karya, Olympos Mt., 25.vi.2014, 1 spec., L. Ernest leg. et coll.; Greece, Thessaly, Larisa pref., Ossa Oros Mt., 5.6.2011, 2 spec., J. Krátký leg. et coll.; Greece, Thessaly, Larisa pref., Ossa Oros Mt., 1.vi.2006, 1 spec., V. Šilha leg. et coll.

***Lasioederma peloponnensis* Toskina, 2013**

*Lasioederma peloponnensis* Toskina, 2013: Toskina, 2013: 19.

**Distribution.** Greece.

***Lasioderma redtenbacheri redtenbacheri* (Bach, 1852)**

*Lasioderma redtenbacheri redtenbacheri* (Bach, 1852): Oertzen, 1886: 254; Jakobson, 1913: 812; Horion, 1961: 241; Crovetti, 1964: 799; Buchelos, 1989: 9; Zahradník, 2007a: 358.

**Distribution.** Central and southern Europe, northern Africa, Middle East, Caucasus, Central Asia, Siberia.

**Material examined.** Greece, Central Macedonia, Thessaloniki pref., Rentina - Modi, 3.-20.vii.2002, 3 spec., Z. Kraus leg. et coll.; Greece, Crete, Heraklio pref., Panasos, Aghia Varvara, 26.iv.2013, 1 spec., S. Benedikt leg., P. Zahradník coll.; Greece, Crete, Lasithi pref., Gournia, Agios Nikolaos env., 8.-9.vi.1992, 1 spec., J. Batelka leg., P. Zahradník coll.

***Lasioderma rhodiicum* Toskina, 2011**

*Lasioderma rhodiicum* Toskina, 2011: Toskina 2011: 56.

**Distribution.** Greece.

***Lasioderma serricorne* (Fabricius, 1792)**

*Lasioderma serricorne* (Fabricius, 1792): Sahlberg, 1913: 17; Jakobson, 1913: 812; Howe, 1958: 14; Pelekassis, 1962: 67; Mourikis & Vassilaina-Alexopoulou, 1975: 148; Buchelos, 1980: 157; Buchelos, 1985: 221; Levinson & Buchelos, 1988: 202; Buchelos, 1989: 9; Buchelos & Athanassiou, 1998: 131; Zahradník, 2007a: 358.

*Lasioderma testaceum* (Duftschmid, 1825): Oertzen, 1886: 255;

**Distribution.** Cosmopolitan, originally obviously from Mediterranean area.

**Material examined.** Greece, Central Macedonia, Thessaloniki, Vrasna, vii.2004, 3 spec., ex larvae from *Prunica* sp., Z. Kraus leg. et coll.; Greece, Thessaly, Trikala pref., Kastraki, vi.2006, 1 spec., ex larvae from *Carpinus* sp., Z. Kraus leg. et coll.; Greece, Peloponnese, Argolida pref., Nafplion, Tolo, vii.2000, 1 spec., L. Ernest leg. et coll.

***Lasioderma torquatum* (Chevrolat, 1859)**

*Lasioderma torquatum* (Chevrolat, 1859): Oertzen, 1886: 255; Yakobson, 1913: 812; Zahradník, 2007a: 358.

**Distribution.** Algeria, France, Greece, Italy, Tunisia.

***Lasioederma tothi* Toskina, 2013**

*Lasioederma tothi* Toskina, 2013: 21.

**Distribution.** Greece.

***Lasioderma triste* Roubal, 1919**

*Lasioderma triste* Roubal, 1919: Roubal, 1919: 38; Español, 1971: 55; Zahradník, 2007a: 358.

**Distribution.** Greece.

**Metholcini Zahradník, 2009*****Metholcus phoenicis* (Fairmaire, 1859)**

*Metholcus cylindricus* (Germar, 1817): Oertzen, 1886: 254; Schilsky, 1898: 71; Yakobson, 1913: 814; Logvinovskiy, 1985: 123; Nardi & Ratti, 1995: 483.

*Metholcus phoenicis* (Fairmaire, 1859): Español, 1992: 123; Zahradník, 2007a: 361.

**Distribution.** Mediterranean.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Stimonía, 20.vi.-4.vii.1992, 1 spec., P. Průdek leg., P. Zahradník coll.; Greece, Central Macedonia, Chalkidiki pref., Vourvouhou, 20.vi.-4.vii.1992, 1 spec., P. Průdek leg., P. Zahradník coll.; Greece, Crete, Chania pref., Agia Marina, 1.vi.2004, 1 spec., J. Januš leg. et coll.; Greece, Crete, Chania pref., Georgioupoli, 26.-30.v.2002, 1 spec., F. Houška leg. et coll.; Greece, Crete, Chania pref., Paleochora, 21.v.-4.vi.2010, 1 spec., P. Kunst leg., R. Mucska coll.; Greece, Greece, Crete, Chania pref., Platanias env., Ildi Oros, 11.vii.2005, 1 spec., J. Simandl leg., P. Zahradník coll.; Crete, Rethymno pref., Panormos, 12.vi.2004, 1 spec., 29.vi.2005, 2 spec., Z. Švec leg. et coll.; Greece, Crete, Rethymno pref., Plakias, 23.-25.vi.2005, 2 spec., Z. Švec leg. et coll.; Greece, Crete, Rethymno pref., Skaleta, 16.vi.2004, 1 spec., Z. Švec leg. et coll.; Greece, Epirus, Preveza pref., Preveza env., Agia Apostoli, Etolla Akarnania, 8.vi.2008, 1 spec., J. Mertlík leg., M. Zýka coll.; Greece, Epirus, Preveza pref., Neochori env., Agia Apostoli, 8.vi.2008, 2 spec., B. Zbuzek leg. et coll.; Greece, Epirus, Preveza pref., Preveza, 15.vi.1999, 3 spec., Z. Švec leg. et coll.; Greece, Epirus, Preveza pref., Preveza, 6.-16.vi.1992, 1 spec., T. Růžicka leg., P. Zahradník coll.; Greece, Epirus, Preveza pref., Preveza, St. Thomas, 12.-14.vii.1997, 2 spec., Z. Černý leg., P. Zahradník et J. Háva coll.; Greece, Ionian Islands, Korfu pref., Benitses env., 23.-26.vi.2003, 1 spec., K. Orszulik leg. et coll.; Greece, Western Greece, Achaia pref., Kalo Rodini, Patra env., 16.vi.2003, 4 spec., S. Benedikt leg., V. Benedikt coll.

***Metholcus rotundicollis* Schilsky, 1898**

*Metholcus rotundicollis* Schilsky, 1898: Reitter, 1901: 27; Heyden, Reitter & Weise, 1906: 430; Pic, 1912b: 41; Yakobson, 1913: 814; Schaufuss, 1916: 726; Español, 1972: 67; Zahradník, 1998: 278; Zahradník et al., 2000: 314; Zahradník, 2007a: 362.

**Distribution.** Middle East, Egypt, Greece, Turkey.

**Material examined.** Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.

**Xyletinini Gistel, 1848*****Pseudoptilinus fissicollis* (Reitter in Saulcy, Reitter et Weise, 1877)**



**Distribution.** Central Europe, Central Asia, Sweden.

**Material examined.** Greece, Thessaly, Larisa pref., Stomio, Ossa Oros Mt., 4.-6.vi.2013, 1 spec., T. Sitek leg. et coll.

**New species for Greece.**

### ***Xyletinus (Calypterus) bucephalus bucephalus* (Illiger, 1807)**

*Xyletinus (Calypterus) bucephalus bucephalus* (Illiger, 1807): Kiesenwetter, 1877: 144; Schilsky, 1898: 64; Pic, 1912b: 48; Jakobson, 1913: 813; Emets & Logvinovskiy, 1977: 415; Español, 1979b: 32; Logvinovskiy, 1985: 112; Zahradník, 2007a: 359.

**Distribution.** Mediterranean area, Arabian Peninsula, Iraq, Portugal.

**Material examined.** Greece, Peloponnese, Korynthia pref., Stylia, 22.v.2007, 1 spec., L. Zombori leg., HNHM coll.

### ***Xyletinus (Calypterus) bucephalus theanus* Reitter, 1901**

*Xyletinus (Calypterus) bucephalus theanus* Reitter, 1901: Gottwald, 1977: 161; Zahradník, 2007a: 359.

*Xyletinus theanus* Reitter, 1901: Reitter, 1901: 29; Heyden, Reitter & Weise, 1906: 431; Roubal, 1942: 20;

*Xyletinus brunnescens* Obenberger, 1917: Obenberger, 1917: 36; Roubal, 1942: 22;

*Xyletinus megatomoides* Roubal, 1917: Roubal, 1917b: 11; Roubal, 1942: 21;

**Distribution.** Bulgaria, Greece, Israel, Egypt, Syria, Turkey.

**Material examined.** Greece, South Aegean islands, Kos pref., Tigaki, 4.-8.ix.1981, 1 spec., T. Palm leg., P. Zahradník coll.; Greece, South Aegean islands, Paros pref., Drios, vii.1997, 2 spec., ex larvae from sheep dung, 2 spec., B. Büche leg. et coll.

**Note:** According to Gottwald (1977) only this subspecies lives in Greece; nominate subspecies living in west mediterran.

### ***Xyletinus (Calypterus) cretus* Toskina, 2007**

*Xyletinus (Calypterus) cretus* Toskina, 2007a: Toskina, 2007a: 78.

**Distribution.** Greece (Crete).

### ***Xyletinus (Calypterus) productus* Toskina, 2006**

*Xyletinus (Calypterus) productus* Toskina, 2006: Toskina, 2006: 227.

**Distribution.** Greece.

### ***Xyletinus (Calypterus) rabusculus* Toskina, 2007**

*Xyletinus (Calypterus) rabusculus* Toskina, 2007a: Toskina, 2007a: 99.

**Distribution.** Greece.

### ***Xyletinus (Calypterus) tenebricus* Toskina, 2007**

*Xyletinus (Calypterus) tenebricus* Toskina, 2007a: Toskina, 2007a: 107.

**Distribution.** Greece.

### ***Xyletinus (Calypterus) tothi* Toskina, 2007**

*Xyletinus (Calypterus) tothi* Toskina, 2007a: Toskina, 2007a: 110.

**Distribution.** Greece.

***Xyletinus (Xyletinus) balcanicus* Gottwald, 1977**

*Xyletinus (Xyletinus) balcanicus* Gottwald, 1977: Gottwald, 1977: 168; Zahradník, 2007a: 360.

**Distribution.** Balcan Peninsula, Italy, Turkey.

**Material examined.** Greece, Epirus, Ioannina, Monodendri, Vicos Mt., 18-19.vi.2010, 1 spec., J. Plecháč leg., P. Zahradník coll.; Greece, Peloponnese, Lakonia pref., Taygetos, 25.v.1990, 1 spec., M. Janata leg., P. Zahradník coll.; Greece, Thessaly, Larisa pref., Spilia, Ossa Oros Mt., 1-2.vi.2007, 1 spec., T. Sitek leg., P. Zahradník coll.; Greece, Thessaly, Trikala pref., Athamania, Tsoumerka Mt., 29.vi.2005, 1 spec., S. Benedikt leg., P. Zahradník coll.; Greece, Western Macedonia, Grevena pref., Prionia - Vermion, 21.v.1937, 1 spec., Bartoň leg., NMP coll. (paratype).

***Xyletinus (Xyletinus) cylindricus* Kofler, 1970**

**Distribution.** Cyprus, Turkey.

**Material examined.** Greece, Crete, Heraklio pref., Pitsidia env., 22.iv.2013, 1 spec., S. Benedikt leg., P. Zahradník coll.; Greece, Crete, Heraklio pref., Stemes env., Kofinas Mt., 24.iv.2013, 1 spec., S. Benedikt leg., V. Benedikt coll. **New species for Greece.**

***Xyletinus (Xyletinus) kofleri* Gottwald, 1977**

*Xyletinus (Xyletinus) kofleri* Gottwald, 1977: Gottwald, 1977: 175; Toskina, 2007b: 86; Zahradník, 2007a: 360.

**Distribution.** Greece.

**Material examined.** Greece, Central Greece, Fthiotida pref., Parnassus Mt., 24.vi.1991, 1 spec., Probst leg., P. Zahradník coll.; Greece, Crete, Chania pref., Omalos, 21.v.2004, 2 spec., J. Januš leg., J. Januš et P. Zahradník coll.; Greece, Crete, Chania pref., Omalos, 22-25.v.1990, 1 spec., Kudrna leg., I. N. Toskina coll.; Greece, Crete, Chania pref., Omalos, 30.v.2004, 1 spec., J. Januš leg., P. Zahradník coll.; Greece, Crete, Lasithi pref., Vai, 16.v.1981, 1 spec., J. Kratochvíl leg., P. Zahradník coll.; Greece, Crete, Rethymno pref., Ideon cave plain, Idi Mt., 10-11.vi.1992, 1 spec., J. Batelka leg., P. Zahradník coll.; Greece, Peloponnese, Lakonia pref., Gonyari, Taygetos Mt., 11-12.vi.1974, 1 spec., J. Horák & V. Švihla leg., USNM coll.; Greece, South Aegean islands, Cyclades pref., Paros I., Naoussa, vii.1997, 1 spec., ex larvae from goat dung, B. Büche leg. et coll.; Greece, Western Greece, Achaea pref., Diakofto, 5.v.2002, 1 spec., J. Macek leg., NMP coll.

***Xyletinus (Xyletinus) konvickai* sp. nov.**

(Figs. 8-10)

**Type material.** Holotype (male): „GRC [Greece] Peloponnese, 37°57'10.99" S 22°25'37.49"V, Ano Trikala, [Peloponnese, Korinthia pref.], 17.5.2009, lgt. O. Konvička", P. Zahradník coll. Paratype: (1 male): „Greece or, Macedonia, Pangeos mts., alpina zone, Ski Resort, 1620 m, Tomáš Sitek lgt.", P. Zahradník coll.

**Description.** Elongate, transversally convex, body length 4.0 mm, the greatest width 2.0 mm. Ratio length : width of elytra 1.7. Black, pubescence grey, short, dense, recumbent. Antennae, palpi and legs black, too.

Head flat, behind clypeus with transversal depression. Front 4 times wider than width of eyes in dorsal view. Their surface shining, with double punctuation – the first fine, dense, puncture almost touched, the second coarse, sparse, distance between punctures twice shorter than their diameter. Pubescence short, inclined forwards. Eyes small, very slightly convex, glabrous, from dorsal view longer than wide. Antennae consist of eleven antennomeres,, serrate, without antennal club (Fig. 8). The first antennomere robust, 1.5 times longer than wide, slightly longer than the second and the third. The second the smallest, slightly triangular and transverse. The third slightly longer than wide. The fourth to eighth transverse, approximately twice wider than long. The ninth and tenth longer than wide. The last antennomere elliptical, twice longer than wide. The last palpomere slightly club-shaped.

Pronotum strongly transverse, twice wider than long (Fig. 9). The greatest width before its base. Posterior angles widely rounded, base of pronotum bent. Surface of pronotum shining, with double punctuation – the first fine, very dense, punctures almost touched, the second coarser and sparser, distance between punctures the same as their diameter. Pubescence short, inclined backwards, on the sides with semierect black hairs. Scutellum semicircular, slightly transverse.

Elytra elliptic, narrowed backward, the greatest width on base of pronotum, without distinct shoulders. Each elytron with twelve narrow striae, but the first and the twelfth very short, extending only one third elytra length. Others extending almost to end of elytra, where connected – the second with the eleventh, the third with the tenth, the fourth with the fifth the sixth with the ninth, and the seventh with the eighth. Surface of elytra shining, with coarse, dense punctures, punctures almost touched. Pubescence short, sparse, inclined backwards.

Legs robust, short, tibiae 1.4 times longer than tarsi. The first tarsomere of posterior leg slightly longer than the second. The third and the fourth short, of the same length, together as long as the second. The fourth deeply heart-shapedly emarginate. The fifth tarsomere to end widened. Claws small, without teeth.

Metasternum without longitudinal ridge. All abdominal ventrites of the same length. Their surface shining, densely and coarsely punctate, punctures almost touched. Pubescence short, dense, inclined backwards.

Projection of genital stirrup ling and narrow (Fig. 10).

Female: Unknown.

**Variability.** Body length 4.1 mm.

**Differential diagnosis.** From similar species from subgenus *Xyletinus* with semierect hairs on sides of pronotum differs by narrowed body and long projection of genital stirrup (*X. sveci* sp. nov.), by black legs, larger body, more transversal pronotum and long projection of genital stirrup (*X. wevalkai*) and by larger body, more transversal pronotum and long projection of genital stirrup (*X. subrotundatus*).

**Name derivation.** Dedicated to the collector of the holotype, Ondřej Konvička, well-known entomologist and specialist in the family Melandryidae.



Figs. 8-10. *Xyletinus (Xyletinus) konvickai* sp. nov.: 8-antenna of male; 9-pronotum; 10-genital stirrup.

***Xyletinus (Xyletinus) laticollis* (Duftschmid, 1825)**

*Xyletinus (Xyletinus) laticollis* (Duftschmid, 1825): Kiesenwetter, 1859: 162; Oertzen, 1886: 254; Schilsky, 1899: 12; Pic, 1912b: 49; Yakobson, 1913: 813; Horion, 1961: 237; Gottwald, 1977: 175; Gottwald, 1982: 113; Zahradník, 2007a: 360.

*Xyletinus (Xyletinus) laticollis robustus* Kofler, 1971: Kofler, 1971: 69.

*Xyletinus (Xyletinus) flavipes* (Castelnau, 1840): Oertzen, 1886: 254; Yakobson, 1913: 813.

*Xyletinus (Xyletinus) fulvicollis* (Reitter, 1890): Schilsky, 1899: 13; Reitter, 1901: 32; Heyden, Reitter & Weise, 1906: 431; Schaufuss, 1916: 726; Español, 1977: 10.

*Xyletinus flavipes fulvicollis* Reitter, 1890: Reitter, 1890: 196.

**Distribution.** Europe, northern Africa, Central Asia, Middle East, Far East, northern Asia.

**Material examined.** Greece, Central Greece, Fokida pref., Sikes env., Giona Mt., 24.vi.2005, 1 spec., S. Benedikt leg., V. Benedikt coll.; Greece, Central Greece, Fithiotida prev., Amfiklia, Parnassus Mt., 21.vi.2005, 2 spec., S. Benedikt leg., V. Benedikt et P. Zahradník coll.; Greece, Central Macedonia, Chalkidiki pref., Sarti env., 1.-12.vi.2010, 1 spec., J. Habarta leg. et coll.; Greece, Central Macedonia, Pieria pref., Litochoro, 10.vi.1999, 1 spec., Z. Švec leg. et coll.; Greece, Central Macedonia, Pieria pref., Nei Pori, 8.-10.vi.1998, 1 spec., J. Simandl leg., P. Zahradník coll.; Greece, Central Macedonia, Pieria pref., Panteleimonas, Kato olympos Mt., 13.-21.vi.1993, 1 spec., J. Růžička leg., P. Zahradník coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.v.2004, 1 spec., Z. Kraus leg. et coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, 3.-20.vii.2002, 2 spec., Z. Kraus leg. et coll.; Greece, Crete, Rethymno pref., Kourna lake, 23.v.2002, 1 spec., J. Simandl leg. et coll.; Greece, Eastern Macedonia and Thrace, Drama pref., Panorama env., Falakro Mt., 3.vi.2008, 1 spec., J. Mertlík leg., M. Zýka coll.; Greece, Eastern Macedonia and Thrace, Kavala pref., Klisma env., Thassos, 22.vi.2006, 1 spec., K. Schön leg., P. Zahradník coll.; Greece, Epirus, Ioannina pref., Leptokaria env., 19.-27.v.1995, 1 spec., Z. Švec leg. et coll.; Greece, Epirus, Ioannina pref., Micro Papigko, 13.vi.1999, 1 spec., Z. Švec leg. et coll.; Greece, Ionian Islands, Zakynthos pref., Argasi env., 5.-14.vi.2013, 1 spec., T. Staněk leg. et coll.; Greece, Northern Aegean islands, Samos pref., Kastania env., 16.vi.2007, 1 spec., K. Schön leg., P. Zahradník coll.; Greece, Peloponnese, Messenia pref., Pylos env., 24.iv.2008, 1 spec., S. Benedikt leg., V. Benedikt coll.; Greece, Thessaly, Larisa pref., Karya, Olympos Mt., 25.6.2014, 1 spec., L. Ernest leg. et coll.; Greece, Thessaly, Larisa pref., Spifa, Ossa Oros Mt., 1.-2.vi.2007, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Larisa pref., Stomio, Spilia, Ossa Oros Mt., 11.vi.2008, 1 spec., J. Mertlík leg., M. Zýka coll.; Greece, Western Greece, Achaia pref., Metochi, Pátra, 14.v.2007, 1 spec., K. Rébl leg. et coll.; Greece, Western Macedonia, Kastoria pref., Eptahori, 15.vi.2010, 1 spec., J. Mertlík leg., M. Zýka coll.

***Xyletinus (Xyletinus) longitarsis longitarsis* Jansson, 1942**

**Distribution.** Europe, Kazakhstan, Mongolia, Turkey.

**Material examined.** Greece, Crete, Chania prov., Omalos, 25.-26.v.2002, 1 spec., F. Houška leg., P. Zahradník coll.

**New species for Greece.**

***Xyletinus (Xyletinus) macedonicus* Toskina, 2007**

*Xyletinus (Xyletinus) macedonicus* Toskina, 2007: Toskina, 2007b: 66.

**Distribution.** Greece.

***Xyletinus (Xyletinus) montanus* Toskina, 2006**

*Xyletinus (Xyletinus) montanus* Toskina, 2006: Toskina, 2006: 248; Toskina, 2007b: 88.

**Distribution.** Greece.

***Xyletinus (Xyletinus) nardi* Zahradník, 2007**

*Xyletinus (Xyletinus) nardi* Zahradník, 2007a: Zahradník, 2007a: 360.

**Distribution.** Azerbaijan, Greece.

**Material examined.** Greece, Thessaly, Larisa pref. Olympiada, 6.vi.1999, 3 spec., Z. Švec leg., Z. Švec et P. Zahradník coll.

***Xyletinus (Xyletinus) ornatus* Germar, 1842**

*Xyletinus (Xyletinus) ornatus* Germar, 1842: Zahradník, 2007a: 360.

**Distribution.** France, Greece, Hungaria, Russia (European area except north part), Siberia, Kazakhstan, Mongolia, Turkey.

***Xyletinus (Xyletinus) pectinatus pectinatus* (Fabricius, 1792)**

*Xyletinus (Xyletinus) pectinatus pectinatus* (Fabricius, 1792): Oertzen, 1886: 254; Yakobson, 1913: 813; Horion, 1961: 238; Zahradník, 2007a: 360.

**Distribution.** Europe, Siberia, Far East.

***Xyletinus (Xyletinus) ruficollis* Gebler, 1833**

*Xyletinus (Xyletinus) ruficollis* Gebler, 1833: Stein & Weise, 1877: 109; Oertzen, 1886: 254; Heyden, Reitter & Weise, 1891: 232; Heyden, Reitter & Weise, 1906: 431; Pic, 1912b: 51; Yakobson, 1913: 813; Schaufuss, 1916: 726; Emets, 1976: 193; Emets & Logvinovskiy, 1977: 418; Logvinovskiy, 1985: 122; Zahradník, 2007a: 361.

**Distribution.** Europe, Siberia, Egypt (Sinai Peninsula), Kazakhstan, Turkey.

***Xyletinus (Xyletinus) sareptanus* Kiesenwetter, 1877**

*Xyletinus (Xyletinus) sareptanus* Kiesenwetter, 1877: Schilsky, 1899: 16; Reitter, 1901: 33; Pic, 1912b: 52; Yakobson, 1913: 813; Schaufuss, 1916: 726; Zaytsev, 1956: 70; Kofler, 1971: 73; Emets, 1976: 193; Zahradník, 2007a: 361.

**Distribution.** Southeastern Europe, Caucasus, Siberia, Central Asia, France, Turkey.

**Material examined.** Greece, Central Greece, Fokia pref., Dafnos env., Ori Vardoussia Mt., 26.v.1997, 2 spec., V. Benedikt leg. et coll.

***Xyletinus (Xyletinus) siteki* sp. nov.**

(Figs. 11-13)

**Type material.** Holotype (male): „Greece bor., Vrasna, [Central Macedonia, Thessaloniki pref.], 26.5.2007. Tomáš Sitek lgt.“, P. Zahradník coll.

**Description.** Elongate, transversally convex, body length 3.5 mm, the greatest width 1.6 mm. Ratio length : width of elytra 1.5. Black, pubescence grey, short, dense, recumbent. Antennae, mandibles, palpi and legs red, antennomeres 4-11 darker, brown.

Head flat, trapezoidal, the widest posteriorly, on sides in middle with short blunt projection to eye. Front 5.2 times wider than width of eye in dorsal view. Their surface shining, with double punctuation - the first fine, very dense, punctures almost touched, the second coarse and sparser, distance between punctures twice as shorter as their diameter. Pubescence short, inclined forwards. Eyes small, slightly convex, longitudinally oval, anteriorly almost straightly severed, glabrous. Antennae consist of eleven antennomeres, serrate, without antennal club (Fig. 11). The first antennomere robust, 2.1 longer than wide, the second the smallest as long as wide, slightly serrate. Antennomeres 3-10 clearly serrate. The third 1.1 longer than wide, antennomere 4-8 slightly transverse, each following more and more, but very similar. The ninth 1.3 times longer than wide, the tenth 1.6 times longer than wide. The last antennomere non-serrate, longly oval, 3 times longer than wide.

Pronotum transverse, twice shorter as wide, bell-shaped (Fig. 12). The greatest wide closely

before their base. Base of pronotum bent. Surface of pronotum shining, with double punctuation - the first fine, very dense, puncture almost touched, the second coarser and sparser, distance between puncture the same as their diameter. Pubescence short, inclined backwards. Scutellum semicircular, slightly transverse.

Elytra oval, before one half slightly narrowed, the greatest width shortly after half; with distinct shoulders. Each elytron with twelve deep and narrow striae, but the first and twelfth very short, extending only to one third length of elytra. Other extending almost to end of elytra, where connected - the second with the eleventh, the third with the tenth, the fourth with the fifth, the sixth with the ninth and with the third and the seventh with the eighth. Surface of elytra shining, with fine dense punctures, punctures almost touched. Pubescence short, sparse, inclined backwards.

Legs robust and short, tibiae with two sharp delayed edges, the greatest width before connected with tarsi, with two small thorns, 1.3 times longer than tarsi. The first tarsomere the longest, 1.3 times longer than the second, this of the same length as the third and the fourth. The third 1.2 times longer than the fourth. The fifth of the same length as the third, but narrower. Claws small, without teeth.


Metasternum without longitudinal ridge. The third and the fourth ventrites slightly narrower than other, the fifth slightly widest. Suture of the first and the second slightly bent in middle. Their surface shining, finely and densely punctuated, punctures almost touched. Pubescence long, fine, dense, inclined backwards.

Projection of genital stirrup long, curved with sharp end (Fig. 13).

Female: Unknown.

**Differential diagnosis.** Shape of pronotum similar as *X. (X.) ater*, but metasternum without longitudinal ridge. Narrowing of elytra before half their length, not present in other *Xyletinus* sensu stricto, is characteristic. From habitually similar *X. (X.) pectinatus* the new species differs by long curved projection on genital stirrup and darker antennae. From similar *X. (X.) longitarsis*, which have also darker antennae and long projection of genital stirrup, it differs by shape of pronotum.

**Name derivation.** Dedicated to the collector of the holotype, Tomáš Sitek, well-known entomologist and specialist in the family Elateridae.



Figs. 11-13. *Xyletinus (Xyletinus) siteki* sp. nov.: 11-antenna of male; 12-pronotum; 13-genital stirrup.

***Xyletinus (Xyletinus) subrotundatus* Lareynie, 1852**

*Xyletinus (Xyletinus) subrotundatus* Lareynie, 1852: Schilsky, 1899: 15; Pic, 1912b: 52; Yakobson, 1913: 813; Horion, 1961: 238; Kofler, 1970; Kofler, 1971: 73; Gottwald, 1977: 176; Emets & Logvinovskiy, 1977: 418; Gottwald, 1982: 114; Logvinovskiy, 1985: 123; Español, 1992: 121; Zahradník, 2007a: 361.

**Distribution.** Central and southern Europe, Middle East, Finland, Kazakhstan, Sweden.

**Material examined.** Greece, Central Macedonia, Pieria pref., Nei Pori, 8.-10.vi.1998, 2 spec., J. Simandl leg., J. Simandl et P. Zahradník coll.; Greece, Central Macedonia, Serres pref., Ampfipolis, 1.-19.viii.2006, 1 spec., Z. Kraus leg. et coll.; Greece, Central Macedonia, Thessaloniki pref., Asprovalta, Vhalkidiki, 15.v.2009, 1 spec., A. Kotán, A. Márkus & T. Németh leg., HHNM coll.; Greece, Central Macedonia, Thessaloniki pref., Vrasna, 8.v.2004, 2 spec., Z. Kraus leg. et coll.; Greece, Epirus, Ioannina pref., Paleochori, Pindos Mt., 14.vi.2010, 1 spec., Z. Malinka leg. et coll.; Greece, Epirus, Ioannina pref., Theodoriana env., Pindos Mt., 27.v.2007, 1 spec., K. Rébl leg. et coll.; Greece, Thessaly, Larisa pref., Omolio, Ossa Oros Mt., 7.vi.1998, 1 spec., J. Simandl leg., USNM coll.; Greece, Thessaly, Larisa pref., Spilia, Ossa Oros Mt., 1.-2.vi.2007, 1 spec., T. Sitek leg. et coll.; Greece, Thessaly, Larisa pref., Stavros refuge, Olympos Mt., 2.vi.2011, 2 spec., O. Konvička leg., O. Konvička et P. Zahradník coll.; Greece, Thessaly, Pieria pref., Panteleimon, Kato Olympos Mt., 13.-21.vi.1993, 1 spec., J. Růžička leg., P. Zahradník coll.; Greece, Thessaly, Trikala pref., Kastraki env., 30.v.2010, 1 spec., A. Trmal leg. et coll.; Greece, Western Macedonia, Grevena pref., Grevena env., Anixi, 31.v.2011, 1 spec., O. Konvička leg. et coll.; Greece, Western Macedonia, Kozani pref., Vathylakkos, 4.v.2004, 1 spec., Z. Kraus leg. et coll.;

***Xyletinus (Xyletinus) sveci* sp. nov.**

(Figs. 14-16)

**Type material.** Holotype (male): „GR [Greece]-Crete, Arkadi, [Heraklio pref.], 8.vi.2004, [Z.] Švec leg., P. Zahradník coll. Allotype (female): „GR [Greece]-KRETE, Amiras [Crete, Heraklio pref.]; 3.6.-5.6.2003, lgt. [V.] Němec“, P. Zahradník coll. Paratypes: (1 male): „Greece bor.-or. Macedonia, [Central Macedonia, Thessaloniki pref.], Olimpiada, Stavros env., 9.vi.2014 50 m, Tomáš Sitek lgt.“ T. Sitek coll.; (1 male) „GR [Greece]-Kefallonia, [Ionian Islands, Kefallonia pref.], Ag. Dimitrios env., 4.6.1997, Ing. K Schön lgt.“, P. Zahradník coll.; (1 male): GR [Greece]-Samos, [Northern Aegean islands, Samos pref.], Kastania env., 16.6.2007, Ing. K. Schön lgt.“, P. Zahradník coll.; (1 female): „Greece-Crete, [Chania pref.], Komitades [35°13'4.77" N, 24°9'57.32" E], 20.5.2007, lgt. O. Konvička“, O. Konvička coll.; (1 female): „Greece, Kreta centr., Ida, [Crete, Rethymno pref.], 1700 m, 4.-12.6.02, 15 km s. Anogia, [damaged, without head], lgt. K. Deneš jun.“, P. Zahradník coll.; (1 male, 1 female): GR [Greece]-Crete, Mts. Ida-Nida [Rethymno pref.], 26.5.2003, lgt. [V.] Němec“, P. Zahradník coll.; (1 male): „I. Jeniš lgt., Omalos Kreta, [Greece, Crete, Chania pref.], 22.-30.5.1990“, P. Zahradník coll.; (2 males): „GR [Greece]-Kreta [Crete], Lassithi prov. [Lasiithi pref.], Selekanos 31.V.-1.VI.04, [V.] Němec lgt.“, P. Zahradník coll.; (1 female): „GR [Greece] Crete bor. Meni, Arkadi env. [Heraklio pref.], 2.6.2013, Švec lgt.“, Z. Švec coll.; (1 male): „Greece-25.5.1997, NE of Vardoússa Mts., DÁFNI vill. env., [Central Greece, Phocis pref.], 1000-1500 m, P. Bulirsch lgt.“, P. Zahradník coll.

**Description.** Shortly elongate, transversally and longitudinally convex, body length 4.0 mm, the greatest width 2.8 mm. Ratio length : width of elytra 1.3. Black, pubescence grey, short, dense, recumbent. Antennae, palpi and legs yellow-red.

Head transversally slightly convex, from dorsal view almost rectangular, clypeus flattened, separated from front by bent rounded edge. Surface of head shining, with two types of punctuation - the first very dense, fine, punctures almost touched, the second sparse, coarse, distance between these punctures are multiples of their diameter. Pubescence recumbent, sparse and long, inclined forwards. Sharp narrow edge going slant to clypeus. Front 4 times wider than width of eye in dorsal view. Eyes oval, slightly convex, almost not protruding from outline of head, glabrous. Antennae consisting of eleven antennomeres, serrate, without antennal club. The first antennomere robust, twice longer than wide, the second small, twice shorter than previous, as long as wide. The third triangular, as long as wide. From fourth to ninth serrate, 1.1 times longer than wide. The tenth also serrate, but 1.2 longer then wide. The last antennomeres elliptical, 2.1 times longer than wide (Fig. 14). The last maxillary palpomere spindleform.

Pronotum transverse (Fig. 15), ratio length : width 0.6. The greatest width in 2/3 before base of pronotum. Anterior angles invisible, posterior angles widely rounded (from dorsal view). Base of pronotum bent. Lateral margin of pronotum with short, dense, black erect hairs. Surface of pronotum shining, with two types of punctuation - the first very dense, fine, punctures almost

touched, the second sparse, coarse and irregular, distance between punctures are multiples of their diameter. Pubescence on disc of pronotum semierect, short, black, inclined backwards. Scutellum trapezoidal, on base of pronotum slightly wider, approximately twice wider than long.

Elytra shortly oval, with distinct shoulders. Each elytron with 11 striae. The first striae, beside suture very short, extending shortly behind scutellum. The second connected with the eleventh before apex of elytra, the third with the tenth, the fourth with the ninth, the fifth with the eighth and the sixth with the seventh, always longly from apex of elytron. Surface of elytra shining, densely and finely punctate, punctures almost touched. Pubescence short, very dense, recumbent, on margin erect.

Legs robust and short, tibiae with two sharp drawn together edges, the greatest width before connected with tarsi, with two small thorns 1.8 times longer than tarsi. The first tarsomere the longest, 1.8 times longer than the second. The second approximately of the same length as the third and fourth. The fourth deeply heart-shapedly emarginate, up to two thirds of length, the fifth tarsomere is inserted in this emargination. The fifth tarsomere twice as long as wide, the greatest width on apex. Claws small, without teeth.

The third and fourth ventrites slightly narrower than others. Their surface shining, very densely and coarsely punctate, punctures almost touched. Pubescence dense, long, inclined backwards.


Projection of genital strirrup short, wide and rounded (Fig. 16).

Female: Antennae less serrate, from fourth to ninth as long as wide.

**Variability.** Body length 4.0-4.1 mm, colour of body reddish (immature specimen?).

**Differential diagnosis.** Only two other *Xyletinus* sensu stricto species have erect hairs on pronotum. *X. (X.) rotundatus* and *X. (X.) wewalkai*. *X. (X.) wewalkai* has only one type of punctuation on pronotum. *X. (X.) subrotundatus* is smaller, narrower, pronotum is enlarged backwards, elytra are matt and apex of genital strirrup is longer and narrower.

**Name derivation.** Dedicated to the collector of the holotype, Zdeněk Švec, well-know entomologist, specialist in the family Leiodidae.



Figs. 14-16. *Xyletinus (Xyletinus) sveci* sp. nov.: 14-antenna of male; 15-pronotum; 16-genital strirrup.



ACKNOWLEDGEMENTS. I would like to thank Jan Batelka (Praha, Czech Republic), Stanislav Benedikt (Plzeň, Czech Republic), Václav Benedikt (Plzeň, Czech Republic), Jean-Claude Besuchet (Genève, Switzerland), Roman Borovec (Sloupno, Czech Republic), Boris Büche (Berlin, Germany), Zdeněk Černý (Mírovce, Czech Republic), Richard Čtvrtečka (Liberec, Czech Republic), Ladislav Daněk (Nymburk, Czech Republic), Ladislav Ernest (Nymburk, Czech Republic), Eduard Ezer (Zlín, Czech Republic), Tomasz Gazurek (Warszawa, Poland), Jan Habarta (Velká nad Veličkou, Czech Republic); Jiří Hájek (Praha, Czech Republic), Jiří Háva (Únětice u Prahy, Czech Republic), Jan Horák (Praha, Czech Republic), Miroslav Janata (Praha, Czech Republic), Jiří Januš (Kladno, Czech Republic), Ivo Jeniš (Lučín, Czech Republic), František Kantner (České Budějovice, Czech Republic), Oldřich Kapler (Světlá, Czech Republic), Ondřej Konvička (Zlín, Czech Republic), Jiří Krátký (Hradec Králové, Czech Republic), A. Kotán (Budapest, Hungary), Zdeněk Kraus (Mikulovice, Czech Republic), Arnošt Kudrna (České Budějovice, Czech Republic), †Jan Macek (Mírošov u Rokycan, Czech Republic), Marion Mantič (Ostrava, Czech Republic), A. Márkus (Budapest, Hungary), Josef Mertlík (Opatovice nad Labem, Czech Republic), Zdeněk Malinka (Opava, Czech Republic), Pavel Moravec (Litoměřice, Czech Republic), V. Němec (Praha, Czech Republic); T. Németh (xxx, Hungary), Kamil Orszulik (Frýdek-Místek, Czech Republic), Jiří Plecháč (Pecka, Czech Republic), R. Plewa (Warszawa, Poland), Pavel Průdek (Brno, Czech Republic), Rolf Röber (Hasselby, Sweden), Jan Růžička (Praha, Czech Republic); Tomáš Růžička (Praha, Czech Republic), Karel Schön (Litvínov, Czech Republic), Jiří Simandl (České Budějovice, Czech Republic), Tomáš Sítek (Ostrava, Czech Republic), Tomáš Staněk (Hradec Králové, Czech Republic), Miroslav Šárovec (České Budějovice), Václav Šilha (Nové Strašecí, Czech Republic), R. Škoda Zdeněk Švec (Praha, Czech Republic), †Vladimír Švihla (Praha, Czech Republic), Antonín Trmal (Sedlčany, Czech Republic), Dušan Vacula (Ostrava, Czech Republic), Jiří Ch. Vávra (Ostrava, Czech Republic), Stanislav Vít (Genève, Switzerland), Pavel Vonička (Liberec, Czech Republic), Bořivoj Zbuzek (Praha, Czech Republic), Lothar Zercher (Berlin, Germany), Miroslav Zýka (Rakovník, Czech Republic) for the kind loan of the material studied or even for giving me their material.

This work was supported by the project of the Ministry of Agriculture of the Czech Republic - Resolution RO0115 (reference number 5774/2015-MZE-17011).

Table 1. List of Greece Ptinidae according to different regions of Greece (mainland and groups of Island; the same structure is used in Fauna Europaea, only Ionian Island are included in addition).

	Family/ Tribe	GR-AEG	GR-CYC	GR-DOD	GR-GRC	GR-ION	GR-KRI
<i>Anitys rubens</i> (Hoffmann, 1803)	Dd				†		
* <i>Anobium hederæ</i> Hssens, 1949	Aa				x		
* <i>Anobium inexpectatum</i> Lohse, 1954	Aa				x		
* <i>Anobium punctatum</i> (DeGeer, 1774)	Aa				x		
* <i>Cacotemnus rufipes</i> (Fabricius, 1792)	Aa				x		
* <i>Caenocara affine</i> (Sturm, 1837)	Dd				x		
<i>Caenocara subglobosum</i> (Mulsant et Rey, 1864)	Dd				x		
<i>Dignomus atticus</i> (Pic, 1902)	PNp			x	x		
<i>Dignomus canaliculatus</i> (Reitter in Brenske et Reitter, 1884)	PNp				x		x
<i>Dignomus frivaldszkyi</i> (Reitter, 1884)	PNp				x		x
<i>Dignomus irroratus</i> (Kiesenwetter, 1851)	PNp		x		x		x
* <i>Dignomus jacqueti</i> (Pic, 1895)	PNp				†		
<i>Dignomus quadricornis</i> (Pic, 1895)	PNp				†		
<i>Dignomus reichei</i> (Boieldieu, 1854)	PNp				x	x	
<i>Dignomus xylopertha</i> (Boieldieu, 1856)	PNp				x		
* <i>Dorcatoma (Dorcatoma) chrysomelina</i> Sturm, 1837	Dd				x		
<i>Dorcatoma (Dorcatoma) palmi</i> Zahradník, 1996	Dd			x			
<i>Dorcatoma (Pilosodorcatoma) setosella setosella</i> Mulsant et Rey, 1864	Dd				x		
<i>Dryophilus densipilis</i> Abeille de Perrin, 1872	DRd			x			
<i>Dryophilus longicollis</i> (Mulsant et Rey, 1853)	DRd						x
<i>Dryophilus pusillus</i> (Gyllenhal, 1808)	DRd				x		
<i>Dryophilus rufescens</i> Pic, 1921	DRd	x					x
<i>Epauloecus unicolor</i> (Piller et Mitterpacher, 1783)	PNp				†		
<i>Episernus graecus</i> sp. nov.	Ee				x		
* <i>Episernus granulatus</i> Weise, 1887	Ee				x		
<i>Episernus taygetanus</i> Mařan, 1941	Ee				x		
<i>Ernobius abietinus</i> (Gyllenhal, 1808)	Ee				x		
<i>Ernobius abietis</i> (Fabricius, 1792)	Ee				x		
<i>Ernobius angusticollis</i> (Ratzeburg, 1837)	Ee				†		
<i>Ernobius cupressi</i> Chobaut, 1899	Ee				†		
<i>Ernobius kailidisi</i> Johnson, 1975	Ee				x		
<i>Ernobius kiesenwetteri</i> Schilsky, 1898	Ee				x		
<i>Ernobius laticollis</i> Pic, 1927	Ee						
* <i>Ernobius lenkae</i> Zahradník, 1996	Ee						x
* <i>Ernobius longicollis</i> (Sturm, 1837)	Ee				x		
<i>Ernobius mollis mollis</i> (Linnaeus, 1758)	Ee			x	x		x
<i>Ernobius nigrinus</i> (Sturm, 1837)	Ee				x		
<i>Ernobius oertzeni</i> Schilsky, 1900	Ee	x			x		
<i>Ernobius pallidipennis</i> Pic, 1902	Ee				x		
<i>Ernobius pini pini</i> (Sturm, 1837)	Ee			x	x		
<i>Ernobius pruinosis</i> (Mulsant et Rey, 1863)	Ee				x		
<i>Ernobius rebli</i> sp. nov.	Ee				x		
* <i>Ernobius reitteri</i> Pic, 1902	Ee			x			
<i>Ernobius robusticornis</i> Mařan, 1941	Ee				x		
<i>Falsogastrallus unistriatus</i> (Zoufal, 1897)	Ag				x		
<i>Gastrallus corsicus</i> Schilsky, 1898	Ag			x	x		x
<i>Gastrallus immarginatus</i> (P.W.J. Müller, 1821)	Ag				x		
* <i>Gastrallus knizeki</i> Zahradník, 1996	Ag				x		
<i>Gastrallus kocheri</i> Español, 1963	Ag						x

<i>Gastrallus laevigatus</i> (Olivier, 1790)	Ag				x	x	x
* <i>Gastrallus mauritanicus</i> Español, 1963	Ag				x		x
* <i>Gastrallus vavrai</i> Zahradnik, 2007	Ag				x		x
<i>Gibbium aequinoctiale</i> Boieldieu, 1854	Gg				x		
<i>Gibbium boieldieui</i> Levrat, 1857	Gg				†		
<i>Gibbium psylloides</i> Czempinski, 1778	Gg				x		
* <i>Hadrobregmus denticollis</i> (Creutzer in Panzer, 1796)	Ah				x		
<i>Hadrobregmus pertinax</i> (Linnaeus, 1758)	Ah				x		
<i>Hedobia pubescens</i> (Olivier, 1790)	EUh				x		
<i>Hemicoelus canaliculatus</i> (C.G. Thomson, 1863)	Aa				x		
<i>Hemicoelus costatus</i> (Aragona, 1830)	Aa				x		
<i>Hemicoelus fulvicornis</i> (Sturm, 1837)	Aa				x		
<i>Hemicoelus rufipennis</i> (Duftschmid, 1825)	Aa				x		
<i>Homophthalmus rugicollis</i> (Mulsant et Rey, 1853)	DRd		x		x		
<i>Hyperisus plumbeum</i> (Illiger, 1801)	Ex		x		x		x
<i>Lasioderma aterrimum</i> Roubal, 1916	XI				x		
<i>Lasioderma baudii</i> Schilsky, 1899	XI				x		x
* <i>Lasioderma bubalus phelipaearu</i> Peyerimhoff, 1926	XI				x		
<i>Lasioderma creatum</i> Toskina, 2013	XI				x		
<i>Lasioderma fuscum</i> (Rey, 1892)	XI				†		
<i>Lasioderma haemorrhoidale</i> (Illiger, 1807)	XI		x		x		x
<i>Lasioderma kiesenwetteri</i> Schilsky, 1899	XI	x			x	x	x
<i>Lasioderma laeve</i> (Illiger, 1807)	XI				x		x
<i>Lasioderma nigellum</i> Toskina, 2013	XI				x		
<i>Lasioderma obscurum</i> (Solsky, 1867)	XI				x	x	x
<i>Lasioderma peloponnensis</i> Toskina, 2011	XI				x		
<i>Lasioderma redtenbacheri redtenbacheri</i> (Bach, 1852)	XI				x		x
<i>Lasioderma rhodiacum</i> Toskina, 2011	XI		x				
<i>Lasioderma serricorne</i> (Fabricius, 1792)	XI				x		
<i>Lasioderma torquatum</i> (Chevrolat, 1859)	XI				x		
<i>Lasioderma tothi</i> Toskina, 2013	XI				x		
<i>Lasioderma triste</i> Roubal, 1919	XI				x		
<i>Mesocoelopus brevistriatus</i> Leiler, 1979	Mm				x		
<i>Mesocoelopus creticus</i> Fairmaire, 1880	Mm		x		x		x
<i>Mesocoelopus niger</i> (P.W.J. Müller, 1821)	Mm	x			x		x
<i>Mesothus ferrugineus</i> (Mulsant et Rey, 1861)	Mm				x		
<i>Metholcus phoenicis</i> (Fairmaire, 1859)	Xm				x	x	x
<i>Metholcus rotundicollis</i> Schilsky, 1898	Xm				x		
<i>Mezium affine</i> Boieldieu, 1856	Gm				x		
<i>Mezium americanum</i> (Castelnau, 1840)	Gm				†		
<i>Nicobium castaneum</i> (Olivier, 1790)	An		x		x		
<i>Niptus</i> ( <i>Niptus</i> ) <i>hololeucus hololeucus</i> (Faldermann, 1835)	PNp				†		
<i>Ochina</i> ( <i>Dulgieris</i> ) <i>latreillii</i> (Bonelli, 1812)	Eo				x		
<i>Ochina</i> ( <i>Ochina</i> ) <i>ferruginea</i> Schilsky 1899	Eo				x		
* <i>Ochina</i> ( <i>Ochina</i> ) <i>hirsuta</i> Seidlitz, 1889	Eo					x	
<i>Ochina</i> ( <i>Ochina</i> ) <i>ptinoides</i> (Marsham, 1802)	Eo				x		
* <i>Oligomerus brunneus</i> (Olivier, 1790)	As				x		
<i>Oligomerus disruptus</i> (Baudi di Selve, 1874)	As				†		
<i>Oligomerus ptilinoides</i> (Wollaston, 1854)	As		x		x		x
<i>Oligomerus retowskii</i> Schilsky, 1898	As				x		
* <i>Priartobium serrifunis</i> Reitter, 1901	As		x				

* <i>Pseudoptinus fissicollis</i> (Reitter in Saulcy, Reitter et Weise, 1877)	Xx				x		
<i>Ptilinus fuscus</i> (Geoffroy in Fourcoy, 1785)	Pp				x		
<i>Ptinomorphus imperialis</i> (Linnaeus, 1767)	EUh				x		
<i>Ptinomorphus magnificus magnificus</i> (Reitter in Leder, 1880)	EUh				x		x
<i>Ptinomorphus perpulchrus</i> (Obenberger, 1917)	EUh				x		
<i>Ptinomorphus regalis</i> (Duftschmid, 1825)	EUh				x		
* <i>Ptinomorphus tatjanae</i> Logvinovskiy, 1978	EUh				x		
<i>Ptinus (Bruchoptinus) brevivittis</i> Reitter, 1881	PNp				x		
<i>Ptinus (Bruchoptinus) femoralis femoralis</i> (Reitter, 1884)	PNp				x		x
<i>Ptinus (Bruchoptinus) henoni</i> Pic, 1897	PNp				x		
<i>Ptinus (Bruchoptinus) pellitus</i> Desbrochers des Loges, 1875	PNp						x
<i>Ptinus (Bruchoptinus) rufipes</i> Olivier, 1790	PNp				x		
<i>Ptinus (bruchoptinus) schatzmayeri</i> Pic, 1934	PNp				x		
<i>Ptinus (Bruchoptinus) torretassoi</i> Pic, 1934	PNp		x		x		
<i>Ptinus (Cyphoderes) bidens</i> Olivier, 1790	PNp				x		
<i>Ptinus (Cyphoderes) raptor</i> Sturm, 1837	PNp				†		
<i>Ptinus (Cyphoderes) schlerethi</i> (Reitter, 1884)	PNp				x		
<i>Ptinus (Gynopterus) crassicornis</i> Kiesenwetter, 1877	PNp				x		
<i>Ptinus (Gynopterus) dubius</i> Sturm, 1837	PNp				x		
<i>Ptinus (Gynopterus) salvatori</i> Pic, 1901	PNp				x		x
<i>Ptinus (Gynopterus) sexpunctatus</i> Panzer, 1789	PNp				†		
<i>Ptinus (Gynopterus) subroseus</i> Reitter, 1888	PNp				x		
<i>Ptinus (Gynopterus) variegatus</i> Rossi, 1794	PNp		x		x		
<i>Ptinus (Pseudoptinus) cumaniensis</i> Pic, 1896	PNp				x		
* <i>Ptinus (Pseudoptinus) nikitani</i> (Brenske et Reitter, 1884)	PNp				x		
<i>Ptinus (Pseudoptinus) oertzeni</i> Reitter, 1888	PNp				x		
<i>Ptinus (Pseudoptinus) rugosicollis</i> Desbrochers des Loges, 1875	PNp				†		
<i>Ptinus (Pseudoptinus) subaeneus</i> (Reitter in Brenske et Reitter, 1884)	PNp				x		x
<i>Ptinus (Ptinus) argolisanus</i> (Reitter, 1884)	PNp				†		
<i>Ptinus (Ptinus) calcaratus</i> Kiesenwetter, 1877	PNp				x		
<i>Ptinus (Ptinus) calcarifer</i> Reitter, 1888	PNp				x		
<i>Ptinus (Ptinus) fur</i> (Linnaeus, 1758)	PNp				x		
<i>Ptinus (Ptinus) gylippus</i> Reitter, 1906	PNp				x		
<i>Ptinus (Ptinus) kiesenwetteri</i> (Reitter in Brenske et Reitter, 1884)	PNp				x		x
<i>Ptinus (Ptinus) kruperi</i> Pic, 1929	PNp				x		
<i>Ptinus (Ptinus) latro</i> Fabricius, 1775	PNp				x		x
<i>Ptinus (Ptinus) obesus</i> Lucas, 1849	PNp				x		
<i>Ptinus (Ptinus) phlomidis</i> Boieldieu, 1854	PNp				x		
<i>Ptinus (Ptinus) pilosus</i> P.W.J. Müller, 1821	PNp		x		x		
<i>Ptinus (Ptinus) rufus</i> Brullé, 1832	PNp				x		
<i>Ptinus (Ptinus) subpilosus</i> Sturm, 1837	PNp			x	x		
<i>Ptinus (Ptinus) tarsalis</i> (Reitter in Brenske et Reitter, 1884)	PNp		x	x	x		x
<i>Ptinus (Ptinus) villiger</i> (Reitter, 1884)	PNp				†		
<i>Ptinus (Tectoptynus) exulans</i> Erichson, 1842	PNp				†		
<i>Ptinus (Tectoptynus) tectus</i> Boieldieu, 1856	PNp						x

<i>Sphaericus (Sphaericus) pinguis</i> (Boieldieu, 1854)	PNs				†		
<i>Stagetus andalusiacus cribricollis</i> (Aubé, 1861)	Dp				†		
* <i>Stagetus byrrhoides</i> (Mulsant et Rey, 1861)	Dp			x			
<i>Stagetus conicicollis</i> (Schilsky, 1899)	Dp				x		
<i>Stagetus curimoides curimoides</i> (Reitter in Brenske et Reitter, 1884)	Dp			x	x	x	
<i>Stagetus curimoides exiguus</i> Roubal, 1917	Dp				x		
<i>Stagetus elongatus</i> (Mulsant et Rey, 1861)	Dp				x	x	x
<i>Stagetus espanoli</i> Toskina 2012	Dp					x	
<i>Stagetus franzi</i> Español, 1969	Dp				x		
<i>Stagetus italicus</i> (Reitter, 1885)	Dp				†		
<i>Stagetus pilula</i> (Aubé, 1861)	Dp				x		
<i>Stagetus propinquus</i> Obenberger, 1917	Dp				x		
<i>Stagetus puncticollis</i> (Brenske et Reitter, 1884)	Dp			x	x		
<i>Stegobium paniceum</i> (Linnaeus, 1758)	As				x	x	x
<i>Trigonogenius globosus</i> Solier, 1849	PNs				†		
<i>Xestobium rufovillosum</i> (DeGeer, 1774)	Ex				x		x
<i>Xyletinus (Calypterus) bucephalus bucephalus</i> (Illiger, 1807)	Xx		x			x	x
<i>Xyletinus (Calypterus) bucephalus theanus</i> Reitter, 1901	Xx	x	x	x	x		x
<i>Xyletinus (Calypterus) cretus</i> Toskina, 2007	Xx						x
<i>Xyletinus (Calypterus) productus</i> Toskina, 2006	Xx				x		
<i>Xyletinus (Calypterus) rabusculus</i> Toskina, 2007	Xx				x		
<i>Xyletinus (Calypterus) tenebricus</i> Toskina, 2007	Xx		x				
<i>Xyletinus (Calypterus) tothi</i> Toskina, 2007	Xx				x		
<i>Xyletinus (Xyletinus) balcanicus</i> Gottwald, 1977	Xx				x		
* <i>Xyletinus (Xyletinus) cylindricus</i> Kofler, 1970	Xx						x
<i>Xyletinus (Xyletinus) kofleri</i> Gottwald, 1977	Xx		x		x		x
<i>Xyletinus (Xyletinus) konvickai</i> sp. nov.	Xx				x		
<i>Xyletinus (Xyletinus) laticollis</i> (Duftschmid, 1825)	Xx			x	x	x	x
* <i>Xyletinus (Xyletinus) longitarsis longitarsis</i> Jansson, 1942	Xx						x
<i>Xyletinus (Xyletinus) macedonicus</i> Toskina, 2007	Xx				x		
<i>Xyletinus (Xyletinus) montanus</i> Toskina, 2006	Xx				x		
<i>Xyletinus (Xyletinus) nardi</i> Zahradnik, 2007	Xx				x		
<i>Xyletinus (Xyletinus) ornatus</i> Germar, 1842	Xx				†		
<i>Xyletinus (Xyletinus) pectinatus pectinatus</i> (Fabricius, 1792)	Xx					x	
<i>Xyletinus (Xyletinus) ruficollis</i> Gebler, 1833	Xx				†		
<i>Xyletinus (Xyletinus) sareptanus</i> Kiesenwetter, 1877	Xx					x	
<i>Xyletinus (Xyletinus) siteki</i> sp. nov.	Xx					x	
<i>Xyletinus (Xyletinus) subrotundatus</i> Lareynie, 1852	Xx					x	
<i>Xyletinus (Xyletinus) sveci</i> sp. nov.	Xx	x				x	x

x - known from this region; † - known from Greece, without specific data for regions; \* - species new to Greece.

GR-AEG - Greece, Northern Aegean Islands; GR-CYC - Greece, Cyclades Islands; GR-DOD - Greece, Dodecanese Islands; GR-GRC - Greece, mainland; GR-ION - Greece, Ionian Islands; GR-KRI - Greece, Crete.

Aa - Anobiinae, Anobiini; Ag - Anobiinae, Gastralini; Ah - Anobiinae, Hadrobregmini; An - Anobiinae, Nicobiini; As - Anobiinae, Stegobiini; Dd - Dorcatominae, Dorcatomini; Dp - Dorcatominae, Prothecini; DRd - Dryophilinae, Dryophilini; Ee - Ernobiinae, Ernobiini; Eo - Ernobiinae, Ochini; Ex - Ernobiinae, Xestobiini; EUh - Eucradinae, Hedobiini; Gg - Gibbiinae, Gibbiini; Gm - Gibbiinae, Meziini; Mm - Mesocoelopodinae, Mesocoelopodini; Pp - Ptilininae, Ptilinini; PNp - Ptininae, Ptinini; PNs - Ptininae, Sphaericini; Xl - Xyletininae, Lasiodermi; Xm - Xyletininae, Metholcini; Xx - Xyletininae, Xyletinini.

## REFERENCES

- BELLÉS X. 1984: *Contribucion al conocimiento de la Familia Ptinidae (Coleoptera). Estudio monografico de la subfamilia Gibbiinae*. Barcelona: Universitat de Barcelona, Centre de Publicacions Interdepartamentals Científic i Extensió Universitària, 40 pp.
- BELLÉS X. 1985: Sistemática, filogenia y biogeografía de la subfamilia Gibbiinae (Coleoptera, Ptinidae). *Treballs del Museu de Zoologia*, No.3: 1-94.
- BELLÉS X. 1996: El género *Dignomus* Wollaston (Coleoptera, Ptinidae). *Bolletí de la Societat d'Historia Natural de Balears* 39: 209-228.
- BELLÉS X. 2002: A synopsis of the subgenus *Cyphoderes* Mulsant & Rey, including the description of *Ptinus* (*Cyphoderes*) *catalonicus* sp. n. (Coleoptera: Ptinidae). *Elytron* 16: 97-105.
- BELLÉS X. & MIFSUD D. 2000: Ptinid beetles (Coleoptera, Ptinidae) from the Maltese islands. *The Central Mediterranean Naturalist* 3: 43-48.
- BOIELDIEU A. A. 1854: [Monographie des Coléoptères de la tribu des Ptinoires.] *Bulletin de la Société Entomologique de France* (3) 2: lxxvii-lxxxv.
- BOROWSKI J. 1993: *Dignomus irroratus* (Kiesenwetter, 1851, (Coleoptera, Ptinidae), nowy dla fauny Polski gatunek chrząszcza, oraz uwagi o jego pozycji systematycznej. *Wiadomości Entomologiczne* 11: 203-205.
- BOROWSKI J. 1996: *Klucz do oznaczania owadów Polski XIX. Chrząszcze - Coleoptera. 42. Pustoszwowe - Ptinidae*. Toruń: Oficyna Wydawnicza Turpress, 45 pp.
- BOROWSKI J. 2002: A world catalogue of the genus *Dignomus* Wollaston (Coleoptera, Ptinidae). *Annals of Warsaw Agricultural University - SGGW, Forestry and Wood Technology*, 52: 35-40.
- BOROWSKI J. 2007: Family Ptinidae Latreille, 1802, subfamilies Gibbiinae Mulsant & Rey, 1868; Ptininae Latreille, 1802. Pp. 328-339. In: LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Volume 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Stenstrup: Apollo Books, 935 pp.
- BRENSKE E. & REITTER E. 1884: Neuer Beitrag zur Käferfauna Griechenlands. Unter Mitwirkung der Herren Dr. Eppelsheim und L. Ganglbauer. *Deutsche Entomologische Zeitschrift* 28: 17-100 + ii pl.
- BUCHELOS C. T. 1980: The population of some Coleoptera species found in storage warehouses of currants and sultanas. *Annales de l'Institut Phytopatologique Benaki (N.S.)* 12: 155-168.
- BUCHELOS C. T. 1985: The Greek Insect fauna of stored products. *Biologia Gallo-Hellenica* 10: 221-227.
- BUCHELOS C. T. 1989: A Contribution to *Lasioderma* spp. and Other Coleoptera Collected from Thistles in Southern Greece. *Entomologica Hellenica* 7: 7-12.
- BUCHELOS C. T. 1994: Coleoptera species, new records from Greek stores. *Annales de l'Institut Phytopatologique Benaki (N.S.)* 17: 101-103.
- BUCHELOS C. T. & ATHANASSIOU C. G. 1998: Beetle species in store-rooms of Central and Southern Greece containing grain, flour, bran and hay: A survey of 44 species. *Annales de l'Institut Phytopatologique Benaki (N.S.)* 18: 129-133.
- BURAKOWSKI B., MROCKZKOWSKI M. & STEFAŃSKA J. 1986: *Katalog fauny Polski. XXIII. Tom 11. Chrząszcze. Dermestoidea, Bostrychoidea, Cleroidea i Lymexyloidea*. Warszawa: Państwowe Wydawnictwo Naukowe, 243 pp. + 1 map.
- CALWER C. G. 1884: *C.G. Calwers Käferbuch. Naturgeschichte der Käfer Europas. Zum Handgebrauche für Sammler. 4. Aufl. Herausgeb. von G. Jäger*. Stuttgart: J. Hoffmann, lvi + 667 pp. + 50 pl.
- CECCHI B. & NARDI G. 1998: Osservazioni su *Ochina latreillii* (Coleoptera, Anobiidae). *Bollettino dell'Associazione Romana di Entomologia* 53: 41-47.
- CHIKATUNOV V. 2000: Catalogue of the beetles (Coleoptera) of Israel and adjacent areas. Tel Aviv: National Collection of Insects, Department of Zoology, The George S. Wise Faculty of Life Sciences, Tel Aviv University, 129 pp.
- CROVETTI A. 1964: Contributi alla conoscenza della entomofauna della *Ferula communis* (L.). IV. Ricerche condotte in Sardegna sugli insetti della parte epigea. *Studi Sarsaresi* 11: 651-908.
- CYMOREK S. 1974: *Familienreihe Teredilia, Kleine Holzwürmer*. In: Schwenke W. (ed.): *Die Forstschädlinge Europas. Ein Handbuch in fünf Bänden. Zweiter Band. Käfer*. Pp. 56-77. Hamburg und Berlin. Verlag Paul Parey, 500 pp.
- DENUX O. & ZAGATTI P. 2010: Coleoptera families other than Cerambycidae, Curculionidae sensu lato, Chrysomelidae sensu lato and Coccinellidae Chapter 8.5. *BioRisk* 4 (1): 315-406. doi:10.3897/biorisk.4.61
- DESBROCHERS DES LOGES J. 1875: *Opuscules entomologiques (Coléoptères). 1<sup>er</sup> Cahier*. Paris: A. Gaudon, 56 pp.
- DOMINIK J. 1966: Obserwacje nad nowo stwierdzonymi w Polsce gatunkami owadów niszczących drewno. *Sylwan* 110 (7): 33-36.
- EMETS V. M. 1976: Obzor vidov zhukov-tochil"shchikov rodov *Xyletinus* Latr. i *Lasioderma* Steph. (Coleoptera, Anobiidae) Mongol'skoy narodnoy respubliki. *Nasekamyye Mongolii* 4: 188-194.
- EMETS V. M. & LOGVINOVSKIY V. D. 1977: Obzor zhukov-tochil"shchikov roda *Xyletinus* Latr. (Coleoptera, Anobiidae) fauny SSSR. *Entomologicheskoe Obozrenie* 56: 409-419.
- ESPAÑOL F. 1963a: Notas sobre Anóbidos VIII. - Los *Gastrallus* mediterráneos. *Publicaciones del Instituto de Biología Aplicada* 35: 5-21.
- ESPAÑOL F. 1963b: Notas sobre anobidos (Coleoptera). XV. Las *Ochina* del Mediterráneo occidental. *Graellsia*

- 20: 287-299.
- ESPAÑOL F. 1964: Los *Xestobium* Motsch. paleárticos (Col. Anobiidae). *Eos, Revista Española de Entomología* 40: 123-137.
- ESPAÑOL F. 1967: Notas sobre anóbidos (Coleoptera). XXX. Contribución al conocimiento del género *Mesocoelopus* Duv. *Publicaciones del Instituto de Biología Aplicada* 43: 85-102.
- ESPAÑOL F. 1969a: Notas sobre Anóbidos (Coleoptera) - XXXV. Los *Stagetus* Woll. del Mediterráneo occidental. *Eos, Revista Española de Entomología* 44: 113-119.
- ESPAÑOL F. 1969b: Notas sobre anóbidos (Col.). XXXVII.-XXXIX. *Publicaciones del Instituto de Biología Aplicada* 46: 49-64.
- ESPAÑOL F. 1970: Notas sobre anóbidos (Col.). XLIX.-L. *Publicaciones del Instituto de Biología Aplicada* 48: 33-48.
- ESPAÑOL F. 1971: Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. No.249 (Col. Anobiidae). *Eos, Revista Española de Entomología* 46: 41-56.
- ESPAÑOL F. 1972: Notas sobre anóbidos - LV. Contribución al conocimiento del género *Metholcus* Duv. *Miscelánea Zoológica* (Barcelona) 3 (2): 61-74.
- ESPAÑOL F. 1977: Sobre algunos Anobiidae (Col.) de Turquía recogidos por el Dr. W. Wittmer (Nota 78). *Mediterranea* 1977 (2): 5-11.
- ESPAÑOL F. 1979a: Generos de Dorcatominae de la fauna Europea (Col., Anobiidae). Nota 92. *Miscelánea Zoológica* (Barcelona) 5: 33-42.
- ESPAÑOL F. 1979b: Contribución al conocimiento de la fauna entomológica del Sudán (Misiones del Prof. H. J. Bremer, 1976-1978). (Col. Anobiidae) (nota 88). *Publicaciones del Departamento de Zoología de la Universidad de Barcelona* 4: 31-37.
- ESPAÑOL F. 1981: Un género y especie nuevos del norte de la India a colocar en la vecindad de *Oligomerus* Redt. (Col., Anobiidae) (Nota 94). *Publicaciones del Departamento de Zoología de la Universidad de Barcelona* 6: 45-55.
- ESPAÑOL F. 1992: *Fauna Iberica. Vol. 2. Coleoptera. Anobiidae*. Madrid. Museo Nacional de Ciencias Naturales. CSIC, 196 pp.
- ESPAÑOL F. 1994: Nuevos datos sobre el género *Falsogastrallus* Pic, 1914 (Coleoptera, Anobiidae). *Zapateri, Revta. Aragon. Ent.* 4: 105-107.
- FAIRMAIRE L. 1880: Descriptions de Coléoptères d'Espagne et de Turquie. *Annales de la Société Entomologique de France* (5) 10: 237-244.
- FALL H. C. 1905: Revision of the Ptinidae of Boreal America. *Transaction of the American Entomological Society* 31: 97-296.
- FAUVEL A. 1889: Rectification au Catalogus Coleopterorum Europae et Caucasi (Suite). *Revue d'Entomologie* 8: 175-203.
- FAUVEL A. 1904: Faune analytique des Coléoptères de la Nouvelle-Caledonie. 2<sup>a</sup> parte. *Revue d'Entomologie* 23: 113-207.
- GEMMINGER M. & HAROLD E. 1869: *Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Tomus VI. Rhipidoceridae, Dascillidae, Malacodermidae, Cleridae, Lymexylonidae, Cupesidae, Ptinidae, Bostrychidae, Cioidae*. Monachii: E.H. Gummi, 1609-1800 + 6 pp.
- GOTTWALD J. 1970: Nové a zajímavé nálezy brouků z Československa (Col.). (4. příspěvek). *Acta Rerum Naturalium Musei Nationalis Slovaci* 16: 119-126.
- GOTTWALD J. 1975: Nové a zajímavé nálezy brouků z Československa (Col.) (6. příspěvek). *Acta Rerum Naturalium Musei Nationalis Slovaci* 21: 143-152.
- GOTTWALD J. 1977: Die paläarktischen *Xyletinus*-Arten (Coleoptera, Anobiidae). *Acta Entomologica Bohemoslovaca* 74: 158-177.
- GOTTWALD J. 1982: Nové a zajímavé nálezy brouků z Československa (Coleoptera) (7. příspěvek). *Entomologické Problémy* 17: 109-115.
- HALPERIN J. & ESPAÑOL F. 1978: The Anobiidae (Coleoptera) of Israel and Near East. *Israel Journal of Entomology* 12: 5-18.
- HEYDEN L. F. J. D. VON, REITTER E. & WEISE J. 1883: *Catalogus Coleopterorum Europae et Caucasi. Editio Tertia*. Berolini, Londini, Parisiis: Libraria Nicolai, Edw. Janson, Lus. Buquet, ii + 238 pp.
- HEYDEN L. F. J. D. VON, REITTER E. & WEISE J. 1891: *Catalogus Coleopterorum Europae, Caucasi et Armeniae rossicae*. Berlin, Mödling Caen. R. Friedländer & Sohn, E. Reitter, *Revue d'Entomologie*, viii + 420 pp.
- HEYDEN L. F. J. D. VON, REITTER E. & WEISE J. 1906: *Catalogus Coleopterorum Europae, Caucasi et Armeniae rossicae. Editio secunda*. Berlin, Paskau, Caen. E. Reitter, 8 unpp.+775 pp.
- HORION A. 1961: *Faunistik der mitteleuropäischen Käfer. Band VIII. Clavicornia 2. Teil. (Thorictidae bis Cistidae). Tereidilia. Coccinellidae*. (Uberlingen, 1961). Genehmigter Nachdruck 1992, Antiquariat Goecke & Evers, xvi + 375 pp.
- HORION A. 1972: Zwölfter Nachtrag zum Verzeichnis der mitteleuropäischen Käfer. *Entomologische Blätter für Biologie und Systematik der Käfer* 68: 9-42.
- HOWE R.W. 1958: A laboratory study of the cigarette beetle, *Lasioderma serricorne* (F.) (Col., Anobiidae) with a critical review of the literature on its biology. *Bulletin of the Entomological Research* 48: 9-56.
- HUBENTHAL W. 1927: Beiträge zur Kenntnis der Gattung *Ernobius* (Col. Anobiidae). *Entomologische Blätter für Biologie und Systematik der Käfer* 23: 82-85.
- IABLOKOFF-KHNZORIAN S. M. & KARAPETYAN A. P. 1991: *Ptinus* - Studien (Coleoptera, Ptinidae). *Entomologische Blätter für Biologie und Systematik der Käfer* 87: 1-65.
- JELÍNEK J. & RÉBL K. 1990: Faunistic record from Czechoslovakia. Coleoptera. *Acta Entomologica Bohemoslovaca* 87:

235-236.

- JOHNSON C. 1975: A Review of the palaearctic Species of the Genus *Ernobius* Thomson (Col., Anobiidae). *Entomologische Blätter für Biologie und Systematik der Käfer* 71: 65-93.
- KARAPETYAN A. P. 1984: Obzor fauny zhukov-tochil"shchikov triby Ernobiini Kavkaza (Coleoptera, Ptinidae). *Biologicheskii Zhurnal Armenii* 37: 947-955.
- KIESENWETTER E. A. H. VON 1859: Beiträge zur Käferfauna Greichlands. Sechstes Stück: Malacodermata, Cleridae, Ptinidae, Anobiidae. *Berliner Entomologische Zeitschrift* 3: 158-191.
- KIESENWETTER E. A. H. VON 1867: Beiträge zur Käferfauna Spaniens. II. Melyridae (Fortsetzung), Ptinidae. *Berliner Entomologische Zeitschrift* 11: 109-134.
- KIESENWETTER E. A. H. VON 1877: *Anobiidae und Cioidae*. Pp. 1-200. In: KIESENWETTER E. A. H. VON & SEIDLITZ G.: Naturgeschichte der Insecten Deutschlands. Erste Abtheilung. Coleoptera 5/1. Berlin. Nicolai, xxviii + 877 pp.
- KOFLER A. 1970: Die paläarktischen *Xyletinus*-Arten des Senckenberg-Museums (Col.: Anobiidae). *Senckenbergiana Biologica* 51: 363-367.
- KOFLER A. 1971: Zur Systematik und Verbeitung europäischer *Xyletinus*-Arten (Col., Anobiidae). *Jahrbuch der Naturwissenschaftlichen Abteilungen am Landesmuseum Joanneum* 1971: 61-74.
- LEILER T.-E. 1979: Beschreibung einer neuen *Mesocoelopus*-Art (Col.,Anobiidae) samt einer Bestimmungstabelle der paläarktischen Arten dieser Gattung. *Entomologische Blätter für Biologie und Systematik der Käfer* 75: 93-97.
- LEILER T.-E. 1985: *Mesocoelopus gambiensis* n.sp., noch eine Anobiidae) Col., aus Asclepiadaceen-Ranken gezüchtet. *Entomologische Blätter für Biologie und Systematik der Käfer* 81: 143-147.
- LEVINSON A. R. & BUCHELOS C. T. 1988: Population dynamics of *Lasioderma serricornis* F. (Col., Anobiidae) in tobacco stores with and without insecticidal treatments: a three year - survey by pheromone and unbaited traps. *Journal of Applied Entomology* 106: 201-211.
- LOGVINOVSKIY V. D. 1977a: K faune i sistematike zhukov-tochil"shchikov podsem. Ernobiinae (Coleoptera, Anobiidae) SSSR. *Entomologicheskoe Obozrenie* 56: 121-131.
- LOGVINOVSKIY V. D. 1977b: Obzor zhukov-tochil"shchikov roda *Lasioderma* Stephens (Coleoptera, Anobiidae) fauny SSSR i Mongolii. *Nasekomyye Mongolii* 5: 278-289.
- LOGVINOVSKIY V. D. 1980: Obzor zhukov-tochil"shchikov roda *Dorcatoma* Herbst (Coleoptera, Anobiidae) fauny SSSR. *Entomologicheskoe Obozrenie* 59: 148-153.
- LOGVINOVSKIY V. D. 1985: *Fauna SSSR. Nasekomyye Zhestkokrylye. Tom XIV, vyp.2. Tochil"shchiki - Semeystvo Anobiidae*. Leningrad. Nauka, 176 pp.
- MARKALAS S. 1991: Rinden- und holzbohrende Insektenarten in Kiefern (*Pinus halapensis*, *P. brutia* und *P. nigra*, nach Waldbränden in Greichenland. *Anzeiger für Schädlingskunde, Pflanzen- und Umweltschutz* 64: 72-75.
- MARSEUL S. A. DE 1886: Catalogue synonymique et géographique des Coléoptères de l'Ancien-Monde Europe et contrées limitrophes en Afrique et en Asie. *L'Abeille* 24: 193-360.
- MAŘAN J. 1941: Nové druhy a formy palearktických Anobiidů. *Sborník Entomologického Oddělení Zemského Musea v Praze* 19: 122-125.
- MOURIKIS M. A. & VASSILAINA-ALEXOPOULOU P. 1975: Report on the most important pests observed on cultivated plants in Greece from 1963 to 1966. *Annales de l'Institute Phytopathologique Benaki (N.S.)* 11: 141-150.
- MULSANT E. & REY C. 1868: *Histoire naturelle des Coléoptères de France. Gibbicolles*. Paris: Deyrolle, Naturaliste, 224 + 16 pp. + xiv pl.
- NARDI G. & RATTI E. 1995: Coleoptera Lyctidae, Bostrichidae e Anobiidae di Pantelleria e Lampedusa. *Naturalista Siciliana, Supplementum* 19: 477-492.
- NOVAK P. 1964: I Coleotteri della Dalmazia. *Atti del Museo Civico di Storia Naturali di Trieste* 24: 53-132.
- OBENBERGER J. 1917: II. Beitrag zur Kenntnis der palaearktischen Käferfauna. *Archiv für Naturgeschichte* A 82 (4): 9-45.
- ORTEZEN E. VON 1886: Verzeichnis der Coleopteren Greichlands und Cretas, nebst einigen Bemerkungen über ihre geographische Verbreitung und 4 die Zeit des Vorkommens einiger Arten betreffenden Sammelberichten. *Berliner Entomologische Zeitschrift* 30: 189-293.
- PELEKASSIS C. E. D. 1962: A catalogue of the more important insects and other animals harmful to the agricultural crops of Greece during the last thirty-year period. *Annales de l'Institute Phytopathologique Benaki (N.S.)* 5: 5-104.
- PIC M. 1896: Descriptions et notes sur divers Coléoptères. *Miscellanea Entomologica* 4: 41-43.
- PIC M. 1901: Description abrégées de Coléoptères Provenant de Grèce. *L'Échange, Revue Linnéenne* 17: 49-52.
- PIC M. 1902a: Nouvelles espèces et variétés de Coléoptères paléarctiques [2<sup>e</sup> article.]. *L'Échange, Revue Linnéenne* 18: 79-80.
- PIC M. 1902b: Nouvelles espèces et variétés de Coléoptères paléarctiques. *L'Échange, Revue Linnéenne* 18 (210): 31-33.
- PIC M. 1903: Espèces et variétés nouvelles de Coléoptères. *L'Échange, Revue Linnéenne* 19: 169-171.
- PIC M. 1908: Descriptions ou diagnoses et notes diverses. Suite. *L'Échange, Revue Linnéenne* 24: 41-46.
- PIC M. 1912a: *Coleopterorum Catalogus. Pars 41 - Ptinidae*. In: JUNK W. & SCHENKLING S. (eds.): *Coleopterorum Catalogus*. Berlin. W. Junk, 46 pp.
- PIC M. 1912b: *Coleopterorum Catalogus. Pars 48 - Anobiidae*. In: JUNK W. & SCHENKLING S. (eds.): *Coleopterorum Catalogus*. Berlin. W. Junk, 92 pp.



- PIC M. 1916: Contribution abrégée. Pour l'étude du genre « *Ernobius* » Thoms. *L'Échange, Revue Linnéenne* 32 (377-378): 1-8 (hors-texte).
- PIC M. 1921a: Notes diverses, descriptions et diagnoses. (Suite). *L'Échange, Revue Linnéenne* 37: 1-4.
- PIC M. 1921b: Notes diverses, descriptions et diagnoses. (Suite). *L'Échange, Revue Linnéenne* 37: 9-10.
- PIC M. 1929: Notes diverses, nouveautés. *L'Échange, Revue Linnéenne* 45: 13-14.
- PIC M. 1934: Notes diverses, nouveautés. (Suite). *L'Échange, Revue Linnéenne* 50: 29-31.
- PORTA A. 1929: *Fauna Coleopterorum Italica. III. Diversicornia*. Piacenza: Stabilimento Tipografico Piacentino, 4 unpp. + 466 pp.
- REITTER E. 1884a: Resultate einer coleopterologischen Sammel-Campagne während den Monaten Februar bis April 1883 auf den jonischen Inseln. *Deutsche Entomologische Zeitschrift* 28: 101-122.
- REITTER E. 1884b: Bestimmungs-Tabellen der europäischen Coleopteren. XI. Bruchidae (Ptinidae). *Verhandlungen des Naturforschenden Vereines in Brünn* 22: 295-323.
- REITTER E. 1888: Neue Coleopteren aus Europa, den angrenzenden Ländern und Sibirien, mit Bemerkungen über bekannte Arten. Fünfter Theil. *Deutsche Entomologische Zeitschrift* 1888: 417-432.
- REITTER E. 1890: Beschreibungen neuer Coleopteren aus Europa, dem Kaukasus, Russisch-Armenien und Turkestan. *Wiener Entomologische Zeitung* 9: 189-198.
- REITTER E. 1901: Analytische Uebersicht der palaeartischen Gattungen und Arten der Coleopteren - Familien. Byrrhidae (Anobiidae) und Cioidae. *Verhandlungen des Naturforschenden Vereines in Brünn* 40: 3-64.
- REITTER E. 1906: Übersicht der Coleopteren-Arten aus der nächsten Verwandtschaft des *Ptinus fur* L., mit weissen Schuppenflecken auf den Flügeldecken. *Wiener Entomologische Zeitung* 25: 281-284.
- REY C. 1892: Remarques en passant. Famille des Térédiles ou Anobides. *L'Échange, Revue Linnéenne* 8: 2-3.
- ROQUES A. 1983: *Les insectes ravageurs des cônes et graines de conifères en France*. Orléans: INRA, 136 pp.
- ROUBAL J. 1917a: Wissenschaftliche Ergebnisse der Bearbeitung von O. Leonhards Sammlungen. *Archiv für Naturgeschichte* A 82 (3): 50-53.
- ROUBAL J. 1917b: Drei neue *Xyletinus*-Formen. *Societas Entomologica* 32: 11.
- ROUBAL J. 1919: Sechs neue palaearktischen Coleopteren. *Archiv für Naturgeschichte* A 83 (7): 36-39.
- ROUBAL J. 1942: *Xyletinus* Latr. (Col. Anob.). Klíč palearktických druhů a nižších systematických jednotek a popisy dvou nových. *Entomologické Listy* 5: 18-25.
- SAALAS U. 1923: Die Fichtenkäfer Finnlands. Studien über die Entwicklungsstadien, Lebensweise und geographische Verbreitung der an *Picea exelsa* Link. lebenden Coleopteren nebst einer Larvenbestimmungstabelle. II. Spezieller Teil 2 und Larve. *Suomalainen Tiedeakatemia Toimituksia* (A) 8: 1-547.
- SAHLBERG J. R. 1903a: Ad cognitionem faunae Coleopterorum Graecae fragmenta. *Öfversigt af Finska Vetenskaps-Societets Förhandlingar* 45 (12): 1-9.
- SAHLBERG J. R. 1903b: Messis hiemalis Coleopterorum Corcyraeorum. Enumeratio Coleopterorum mensibus novembri - februario 1895 - 1896 et 1898 - 1899 nec non primo vere 1896 in Insula Corcyra collectorum. *Öfversigt af Finska Vetenskaps-Societets Förhandlingar* 45 (11): 62-65.
- SAHLBERG J. R. 1913: Messis nova hiemalis Coleopterorum Corcyraeorum. Enumeratio Coleopterorum mensibus Novembri et Decembri 1903 in insula Corcyra collectorum quae ibi antea haud vel rarissime observari. *Öfversigt af Finska Vetenskaps-Societets Förhandlingar* 55 (12): 1-28.
- SEIDLITZ G. 1889: *Fauna Baltica. Die Käfer (Coleoptera, der Deutschen ostseeprovinzen Russlands. Zweite neu bearbeitete Auflage*. Königsberg: Hartung'sche Verlagsdruckerei, 81-128 + 337-512.
- SEIDLITZ G. 1891: *Fauna Transsylvanica. Die Käfer (Coleoptera) Siebenbürgens*. Königsberg: Hartung'sche Verlagsdruckerei, xlix-lvi + 129-192 + 545-914.
- SHARP D. 1916: Additions and corrections in the genus *Ernobius*; with notes on the copula. *The Entomologists Monthly Magazine* 52: 219-224.
- SCHAUFUSS C. 1916: *Calwer's Käferbuch. Einführung in die Kenntnis der Käfer Europas. Sechste Auflage*. Stuttgart: E. Schweizerbart'sche Verlagsbuchhandlung Nägele & Dr. Sprösser, Band II. 390 pp.
- SCHILSKY J. 1898: *Die Käfer Europa's. XXXV*. Nürnberg: von Bauer und Raspe (Emil Küster), viii + 100 pl. [268 pp.] + 43 unpp.
- SCHILSKY J. 1899: *Die Käfer Europa's. XXXVI*. Nürnberg: von Bauer und Raspe (Emil Küster), iv + 100 pl. [259 pp.] + 54 unpp.
- SCHILSKY J. 1900: *Die Käfer Europa's XXXVII*. Nürnberg: von Bauer und Raspe (Emil Küster), iv + 100 pl. [240 pp.] + 22 unpp.
- STEIN J. P. E. F. & WEISE J. 1877: *Catalogi Coleopterorum Europae. Editio secunda*. Berolini, Londini, Parisiis: Nicolai, E. Janson, L. Buquet, 212 pp.
- TOSKINA I. N. 2001: Three new Palaeartic species of the genus *Ptinomorphus* Mulsant & Rey, 1868 (Coleoptera: Anobiidae). *Russian Entomological Journal* 10: 143-151.
- TOSKINA I. N. 2003: The genera *Gastrallus* Jacquelin du Val, 1860, and *Falsogastrallus* Pic, 1914 (Coleoptera: Anobiidae) of Palaeartic Eurasia. *Russian Entomological Journal* 12: 187-197.
- TOSKINA I. N. 2006: New Palaeartic species of the genus *Xyletinus* Latreille, 1809 (Coleoptera: Anobiidae). *Zoosystematica Rossica* 14: 223-260.

- TOSKINA I. N. 2007a: New species of the subgenera *Pseudocalypterus* and *Calypterus* of the genus *Xyletinus* Latreille, 1809, and key to Palaearctic Calypterus (Coleoptera: Anobiidae). *Elytron* 21: 63-120.
- TOSKINA I. N. 2007b: Some new species and key to palaearctic species of the genus *Xyletinus* Latreille, 1809, subgenus *Xyletinus* s. str. (Coleoptera: Anobiidae). *Elytron* 20 [2006]: 55-98.
- TOSKINA I. N. 2011: Dva novykh sredizemnomorskikh vida tochilshchikov podsemeystva Xyletininae (Coleoptera: Ptinidae). *Byulleten Moskovskogo Obshchestva Ispytateley Prirody. Otdel Biologicheskoy Prirody* 116 (6): 54-57.
- TOSKINA I. N. 2012: Neskolko novykh palearkticheskikh vidov roda *Stagetus* Wollaston, 1861 (Coleoptera: Ptinidae: Dorcatominae). *Byulleten Moskovskogo Obshchestva Ispytateley Prirody. Otdel Biologicheskoy Prirody* 117 (2): 16-28.
- TOSKINA I. N. 2013: Novye yuzhnoevropeyskie tochilshchikov roda *Lasioderma* Stephens, 1835 (Coleoptera: Ptinidae: Xyletininae). *Eversmannia* 34: 3-24.
- WHITE R. E. 1990: *Lasioderma haemorrhoidale* (Ill.) now established in California, with biological data on *Lasioderma* species (Coleoptera: Anobiidae). *The Coleopterists Bulletin* 44: 344-348.
- YAKOBSON G. G. 1913: *Zhuki Rossii i zapadnoy Evropy*. St. Peterburg: Izd. A. F. Devriena, 721-864 pp. + 76-83 pl.
- ZAHRADNÍK P. 1996: Four new species of Anobiidae (Coleoptera) from Turkey and Rhodes Island. *Folia Heyrovskyana* 4: 93-101.
- ZAHRADNÍK P. 1998: Anobiidae of Turkey (Coleoptera). *Klapalekiana* 34: 263-286.
- ZAHRADNÍK P. 2000: *Ernobius besucheti* sp.n. (Coleoptera: Anobiidae) from Switzerland. *Folia Heyrovskyana* 8: 133-136.
- ZAHRADNÍK P. 2007a: Family Ptinidae Latreille, 1802 [- Gibbiinae Mulsant & Rey, 1868, Ptininae Latreille, 1802]. Pp. 339-362. In: LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Volume 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Stenstrup: Apollo Books, 935 pp.
- ZAHRADNÍK P. 2007b: Contribution to knowledge of the tribe Gastralini (Coleoptera: Bostrichoidea: Anobiidae) - I. New species of the genus *Gastrallus* from Turkey, with review of the Palaearctic species. *Studies and Reports of District Museum Prague-East. Taxonomical Series* 3: 171-178.
- ZAHRADNÍK P. 2010: Contribution to knowledge of the tribe Gastralini (Coleoptera: Bostrichoidea: Anobiidae) - IV. Review of the genus *Falsogastrallus*, with description of new species. *Studies and reports. Taxonomical Series* 6: 271-275.
- ZAHRADNÍK P., CHIKATUNOV V. & PAVLIČEK T. 2000: Catalogue of beetles (Coleoptera) in Israel and adjacent areas: 2. Anobiidae. *Klapalekiana* 36: 307-319.
- ZAHRADNÍK P. & HÁVA J. 2014: Catalogue of the world genera and subgenera of the superfamilies Derodontoidea and Bostrichoidea (Coleoptera: Derodontiformia, Bostrichiformia). *Zootaxa* 3754 (4): 301-352.
- ZAYTSEV F. A. 1956: Zhuki-Tochil'shchiki v faune Zakavkaz'ya (Coleoptera, Anobiidae). [Anobiid beetle in the fauna of Transcaucasia (Coleoptera, Anobiidae)]. *Trudy Instuta Zoologii AN Gruzinskoy SSR* 14: 49-77.
- ZOUFAL V. 1897: Zwei neue *Gastrallus*-Arten. *Wiener Entomologische Zeitung* 16: 206.